






Supplementary information to:

Original article:

**THE BOVINE DIALYZABLE LEUKOCYTE EXTRACT,
IMMUNEPOTENT CRP, SYNERGICALLY ENHANCES
CYCLOPHOSPHAMIDE-INDUCED BREAST CANCER CELL DEATH,
THROUGH A CASPASE-INDEPENDENT MECHANISM**

Ana Luisa Rivera-Lazarín ^{a,+} , Ana Carolina Martínez-Torres ^{a,+*} ,
Rafael de la Hoz-Camacho ^a , Olga Liliana Guzmán-Aguillón ^a ,
Moisés Armides Franco-Molina ^a , Cristina Rodríguez-Padilla ^{a,b} 

^a Universidad Autónoma de Nuevo León, Facultad de Ciencias Biológicas, Laboratorio de
Inmunología y Virología, Monterrey 66455, México


^b LONGEVEDEN S.A. de C.V.

⁺ Co-first authors


^{*} **Corresponding author:** Ana Carolina Martínez-Torres, Universidad Autónoma de
Nuevo León, Facultad de Ciencias Biológicas, Laboratorio de Inmunología y Virología,
Monterrey 66455, México. E-mail: ana.martinezto@uanl.edu.mx

<https://dx.doi.org/10.17179/excli2022-5389>


This is an Open Access article distributed under the terms of the Creative Commons Attribution License
(<http://creativecommons.org/licenses/by/4.0/>).




Supplementary Figure 1: A) MCF-7, B) MDA-MB-231 and C) 4T1 cells were treated for 24 h. Flow cytometry was used to measure cell death analyzed by AnnV/PI staining. Representative dot plots from at least three independent experiments.



Supplementary Figure 2: Representative dot plots of cell death obtained by AnnV/PI of **A)** MCF-7, **B)** MDA-MB-231 and **C)** 4T1 cells pre-treated in presence or absence of QVD, and then treated with ICRP CC₅₀ or CTX CC₅₀ for 24 h, using flow cytometry.



Supplementary Figure 3: Representative dot plots of cell death obtained by AnnV/PI of **A)** MCF-7, **B)** MDA-MB-231 and **C)** 4T1 cells pre-treated in presence or absence of NAC, and then treated with ICRP CC₅₀ or CTX CC₅₀ for 24 h, using flow cytometry.



Supplementary Figure 4: PBMC obtained from healthy donors were treated with CC₂₀ or CC₅₀ of ICRP or CTX for 24 h and cell death was evaluated by AnnV/PI staining using flow cytometry. **A)** Dot plots are representative of means of triplicates from at least three independent experiments.