

Supplementary information to:

Letter to the editor:

**CYCLOPHOSPHAMIDE-ASSOCIATED PARTIAL PROXIMAL
TUBULAR DYSFUNCTION**

Rutvikkumar Jadvani , Angel Juarez , Chintan V. Shah* 

Division of Nephrology, Hypertension, and Renal Transplantation,
University of Florida College of Medicine, Gainesville, Florida, USA

* **Corresponding author:** Chintan V. Shah, MD, Division of Nephrology, Hypertension,
and Renal Transplantation, University of Florida College of Medicine, 1600 SW Archer
Road, Room CG-98, Gainesville, FL 32610, Office: (352) 294-8790;
E-mail: shahc@ufl.edu

<https://dx.doi.org/10.17179/excli2025-8622>

This is an Open Access article distributed under the terms of the Creative Commons Attribution License
(<https://creativecommons.org/licenses/by/4.0/>).

Table S1: Serum and urine chemistry

	Serum Chemistry	
	On presentation	Sodium Potassium Phosphate Uric acid Magnesium Total serum CO ₂ Creatinine PTH 25-OH, Vitamin D 1-25 dihydroxy vitamin D
	Serum phosphorus level	Fractional excretion of phosphorus (FePO₄)
On presentation	3.3 mg/dL	21 %
At 3 months follow-up ^a	2.7 mg/dL	11 %
At 9 months follow-up ^a	2.4 mg/dL	14.63 %
	Urine Amino Acids	
		Reference Range
At 9 months follow-up	Alanine 710 µmol/g Leucine 53 µmol/g Taurine 10703 µmol/g Tryptophan 101 µmol/g	60-500 µmol/g < 45 µmol/g < 3200 µmol/g 15-95 µmol/g

^a Serum phosphorus levels were measured after holding oral phosphorus supplements for 4 weeks.