

Supplementary information to:

Original article:

UP-REGULATION OF MIR-155 POTENTIATES CD34⁺ CML STEM/PROGENITOR CELLS TO ESCAPE FROM THE GROWTH-INHIBITORY EFFECTS OF TGF-B1 AND BMP SIGNALING

Touba Mahdloo¹ , Pantea Sahami² , Reihaneh Ramezani² , Mojtaba Jafarinia¹ ,
Hamedreza Goudarzi¹ , Sadegh Babashah^{3,*} 

¹ Department of Genetics, Faculty of Basic Sciences, Islamic Azad University, Marvdasht, Iran


² Department of Biomedical Sciences, Women Research Center, University of Alzahra, Tehran, Iran

³ Department of Molecular Genetics, Faculty of Biological Sciences, Tarbiat Modares University, Tehran, Iran

* **Corresponding author:** Sadegh Babashah, Ph.D., Department of Molecular Genetics, Faculty of Biological Sciences, Tarbiat Modares University, P.O. Box: 14115-154, Tehran, Iran, Tel.: +98 21 8288 4468, fax: +98 21 8288 4717, E-mail: sadegh.babashah@gmail.com; babashah@modares.ac.ir

<http://dx.doi.org/10.17179/excli2021-3404>

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>).



Supplementary Figure 1: Fold change of miR-155 in CD34⁺ CML cells following transfection with anti-miR-155. miRNA level was measured by qRT-PCR and normalized to U6 snRNA as internal control.