

Supplementary information to:

ANALYSIS OF NATURAL KILLER CELL FUNCTIONS IN PATIENTS WITH HEREDITARY HEMOCHROMATOSIS

Vivian Bönnemann¹, Maren Claus¹, Barbara Butzeck², Daniela Collette³, Peter Bröde¹, Klaus Golka¹, Carsten Watzl¹

¹ Department for Immunology, Leibniz Research Centre for Working Environment and Human Factors (IfADo) at TU Dortmund, Dortmund, Germany


² Hämochromatose-Vereinigung Deutschland e.V. HVD, European Federation of Associations of Patients with Haemochromatosis EFAPH, Hattingen, Germany

³ Gemeinschaftspraxis für Hämatologie und Onkologie, Dortmund, Germany

* **Corresponding author:** Carsten Watzl, Leibniz Research Centre for Working Environment and Human Factors (IfADo) at TU Dortmund, Ardeystrasse 67, 44139 Dortmund, Germany, E-mail: watzl@ifado.de

<http://dx.doi.org/10.17179/excli2020-1116>

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>).



Supplementary Figure 1: Analysis of leukocyte subpopulations by multicolor flow cytometry.
A) Gating strategy for a general overview of lymphocyte and monocyte subpopulations. **B)** Quantitative analysis of flow cytometry data for HH patients compared to healthy controls (each n=21): Mean \pm SD of **B** % NK cells of lymphocytes, **C** % T cells of lymphocytes, **D** % B cells of lymphocytes, **E** % CD14 positive monocytes of monocytes (in PBMC). The data is analyzed by Wilcoxon rank-sum test, also called Mann-Whitney U test; no statistical significance was detected.