

Supplementary material to:

PREDICTION OF AROMATASE INHIBITORY ACTIVITY USING THE EFFICIENT LINEAR METHOD (ELM)

Watshara Shoombuatong¹, Veda Prachayasittikul^{1,2}, Virapong Prachayasittikul², Chanin Nantasenamat^{1,2,*}

¹ Center of Data Mining and Biomedical Informatics, Faculty of Medical Technology, Mahidol University, Bangkok 10700, Thailand

² Department of Clinical Microbiology and Applied Technology, Faculty of Medical Technology, Mahidol University, Bangkok 10700, Thailand

* Corresponding author: E-mail: chanin.nan@mahidol.ac.th (C.N.); Phone: +66 2 441 4371; Fax: +66 2 441 4380

<http://dx.doi.org/10.17179/excli2015-140>

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>).

Supplementary Table S1: Initial and 10 optimal parameter sets of ELM for predicting steroid AIs

Feature	Initial	1	2	3	4	5	6	7*	8	9	10
C-025	0.13	-0.21	0.08	0.36	0.19	0.19	0.17	0.13	0.11	0.33	-0.06
ESpm13r	6.70	-13.48	-7.93	-15.55	-11.25	-11.25	-9.18	-15.24	-11.07	-9.62	-9.77
ESpm14u	2.95	7.30	7.12	11.57	6.33	6.33	9.18	9.23	9.71	8.05	4.53
ESpm10r	5.84	2.57	-0.90	5.65	4.32	4.32	2.34	5.30	3.99	0.00	5.23
ESpm12x	5.49	-0.54	-2.72	-6.68	-6.27	-6.27	-8.32	-5.94	-3.09	-8.67	-3.46
ESpm15d	0.12	0.13	0.09	0.21	0.12	0.12	0.19	0.15	0.13	0.32	0.26
ESpm10x	6.33	1.01	2.12	3.74	6.76	6.76	3.79	5.43	-2.69	9.73	1.85
H-050	0.07	0.03	0.05	0.06	0.09	0.09	0.05	0.13	0.03	0.07	0.09
nBM	0.04	0.03	0.00	-0.02	-0.03	-0.03	0.03	0.00	0.10	-0.04	0.03
MATS6p	1.52	3.93	3.18	3.88	3.34	3.34	3.84	3.85	2.37	3.08	4.48
MATS6e	1.52	-2.46	-2.44	-2.63	-1.88	-1.88	-3.24	-3.15	-1.80	-2.29	-4.02
GATS6m	0.94	-2.44	-1.92	-2.27	-1.69	-1.69	-1.99	-2.32	-1.48	-2.12	-2.58
GATS6p	1.03	2.62	1.35	2.41	1.89	1.89	1.40	1.58	1.01	1.86	1.72
piPC07	0.29	0.62	0.64	0.74	0.26	0.26	0.72	0.23	0.46	0.02	0.41
GGI1	0.08	0.02	-0.01	-0.02	-0.02	-0.02	-0.06	-0.03	0.03	-0.11	-0.09

* Experiment 7 provided the highest performance for predicting steroid AIs

Supplementary Table S2: Initial and 10 optimal parameter sets of ELM for predicting non-steroidal AIs

Feature	Initial	1	2	3	4*	5	6	7	8	9	10
nAB	0.01	-0.02	-0.03	-0.02	-0.02	-0.02	-0.03	-0.02	-0.02	-0.02	-0.02
piPC05	0.31	2.17	1.94	1.83	2.10	1.81	2.16	2.21	1.92	2.06	1.96
BEHp4	0.37	-1.92	-2.30	-2.23	-2.19	-2.01	-1.97	-1.83	-1.93	-2.18	-1.79
Yindex	0.24	0.88	1.00	0.96	1.08	0.95	0.96	0.96	1.06	1.04	1.09
F04[O-O]	0.02	0.13	0.11	0.12	0.13	0.13	0.11	0.13	0.12	0.12	0.11
EEig09x	0.09	0.25	0.20	0.22	0.23	0.21	0.21	0.22	0.21	0.24	0.19
GATS5e	0.08	0.17	0.17	0.14	0.15	0.12	0.13	0.13	0.13	0.15	0.11
nCp	0.03	0.11	0.10	0.10	0.11	0.11	0.11	0.14	0.10	0.10	0.12
ATS5p	0.16	-0.51	-0.43	-0.41	-0.46	-0.46	-0.48	-0.48	-0.54	-0.49	-0.54
piPC08	0.16	0.90	0.94	1.05	0.95	1.03	0.89	0.98	0.95	1.03	0.90
EEig04x	0.14	-0.37	-0.51	-0.33	-0.40	-0.34	-0.37	-0.35	-0.49	-0.30	-0.35
piPC06	0.31	-2.92	-2.88	-2.97	-3.00	-2.92	-2.85	-3.11	-2.85	-3.02	-2.79
ESpm01x	0.53	1.06	1.88	1.24	1.25	1.22	1.06	1.24	1.68	1.03	1.33
F04[C-O]	0.01	-0.01	-0.01	-0.01	-0.01	-0.01	-0.01	-0.01	-0.01	-0.01	-0.01
BEHe4	0.38	1.89	2.05	2.15	2.14	1.94	2.05	1.66	1.92	2.21	1.73

* Experiment 4 provided the highest performance for predicting non-steroidal AIs