

## Review article:

# MicroRNAs IN ENDOMETRIAL CANCER: RECENT ADVANCES AND POTENTIAL CLINICAL APPLICATIONS

Megumi Yanokura, Kouji Banno\*, Miho Iida, Haruko Irie, Kiyoko Umene, Kenta Masuda, Yusuke Kobayashi, Eiichiro Tominaga, Daisuke Aoki

Department of Obstetrics and Gynecology, Keio University School of Medicine, Tokyo, Japan

\* Correspondence: Dr. Kouji Banno, Department of Obstetrics and Gynecology, School of Medicine, Keio University, Shinanomachi 35 Shinjuku-ku, Tokyo 160-8582, Japan, Phone: +81-3-3353-1211; Fax: +81-3-3226-1667; E-mail: kbanno@z7.keio.jp

<http://dx.doi.org/10.17179/excli2014-590>

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>).

## ABSTRACT

Endometrial cancer is a common malignant gynecological tumor, but there are few biomarkers that are useful for early and accurate diagnosis and few treatments other than surgery. However, use of microRNAs (miRNAs) that induces gene downregulation in cells may permit effective and minimally invasive diagnosis and treatment. In endometrial cancer cells, expression levels of miRNAs including miR-185, miR-210 and miR-423 are upregulated and those of miR-let7e, miR-30c and miR-221 are downregulated compared to normal tissues, and these miRNAs are involved in carcinogenesis, invasion and metastasis. miRNAs with expression changes such as miR-181b, miR-324-3p and miR-518b may be used as prognostic biomarkers and transfection of miR-152 may inhibit cancer growth. However, most current studies of miRNAs are at a basic level and further work is needed to establish clinical applications targeting miRNAs.

**Keywords:** microRNA, endometrial cancer, biomarker, OncomiR, tumor suppressor miR

## INTRODUCTION

microRNA (miRNA) is of interest as a regulatory mechanism in genomic expression. The human genome project identified approximately 20,000 genes in humans, but only 2 % of these genes are translated into proteins (International Human Genome Sequencing Consortium, 2004). A large-scale transcriptome analysis showed that RNAs are also transcribed in several noncoding regions, in which gene sequences of proteins are not encoded. These so-called non-coding RNAs (ncRNAs) have complex functions including regulation of chromatin structure, splicing, and protein silencing.

miRNA is the most important class of ncRNAs. miRNAs are small RNA molecules of about 22 nucleotides that induce gene silencing. About 2,000 miRNAs have been

found in humans and these molecules are important for biological activities including development; cell differentiation, proliferation and death; and metabolism (Stefani and Slack, 2008; Elbashir et al., 2001). Various diseases are caused by breakdown of gene regulatory networks due to changes in miRNA levels, and the relationship of miRNAs with cancer is under investigation.


One miRNA regulates many messenger RNAs (mRNAs), while cancer does not develop due to a single cause, but due to breakdown of multiple mechanisms. Therefore, in contrast to common drugs, miRNAs may be used as treatment through comprehensive regulation of expression of molecules related to cancer. Use of miRNAs may also contribute to elucidation of oncogenic and homeostatic mechanisms. In this article,

we summarize the latest studies on application of miRNAs in diagnosis and treatment for endometrial cancer, based on malignant transformation associated with abnormal miRNA expression.

### miRNA AND CANCER

miRNA-induced RNA silencing is involved in homeostatic maintenance of organisms and breakdown of this mechanism causes disease. Cancer is among the best-studied diseases caused by failure of miRNA regulation. miRNAs are implicated in cancer development and progression, and expression patterns of miRNAs in normal tissues differ from those of cancer tissues. miRNAs involved in cancer are classified into oncogenic miRNAs (oncomiRs), which are upregulated in tumor tissues; and tumor suppressor miRNAs (tumor suppressor miRs), which are downregulated in tumor tissues and inhibit cancer. OncomiRs target mRNAs that suppress tumor growth, whereas tumor suppressor miRs target mRNAs that promote tumor growth. Failed regulation of both types of miRNAs promotes oncogenesis, proliferation and invasion of cancer cells, and the epithelial-mesenchymal transition (EMT) (Figure 1).

Involvement of miRNAs in human cancer was first shown in B cell chronic lymphocytic leukemia (CLL), in which expression of miR-15 or miR-16 was found to decrease (Calin et al., 2002). In cancers, miR-17-92 is upregulated in lung cancer cells (Hayashida et al., 2005). A large-scale analysis of miRNA expression in 540 patients with lung, breast, gastric, prostate, colon and pancreatic cancer showed that expression profiles of miRNAs in cancer tissues differ from those in normal tissues (Volinia et al., 2006). miR-21 was particularly upregulated in all types of cancer, indicating that it is likely to be involved in cancer onset. Subsequently, miR-21 was shown to regulate transcriptional repressors including NFIB (Fujita et al., 2008). Upregulation of miR-21 downregulates transcriptional repressors, resulting



**Figure 1:** Changes in miRNA expression and oncogenic mechanisms. Upregulation of oncomiRs inhibits expression of tumor-suppressor mRNAs, resulting in enhanced oncogenesis because tumor suppression effects are lost. In contrast, downregulation of tumor-suppressor miRs reduces inhibition of expression of oncogene mRNAs, and the resultant upregulated expression of these mRNAs facilitates oncogenesis.

in transcriptional activation; therefore, miR-21 is considered to be a typical oncomiR.

miRNAs that are downregulated in cancer have also been found. For example, let-7, which was found in *C. elegans* and is involved in cell growth, is a typical miRNA precursor that is downregulated in breast, lung and gastric cancer. let-7 targets oncogenes including RAS and high mobility group AT-hook 2 (HMGA2) and has been shown to be a tumor suppressor miR (Johnson et al., 2005; Lee and Dutta, 2007; Yu et al., 2007). Some miRNAs have carcinoma-specific differential expression. The relationship of many of these miRNAs with their target genes is unknown, but miR-19 is necessary for inhibition of apoptosis and oncogenesis in lymphocytes (Olive et al., 2009). Chen et al. (2007) found that undifferentiated cell clusters including embryonic stem (ES) cells, embryoid bodies and mouse 11-day embryos had expression profiles of miRNAs that were simpler than those of mature somatic cells. These results suggest that the differentiation stage may be defined by profiles of specific miRNAs. It is also of note that miRNA profiles in undifferentiated stem cells are similar to those of cancer cells.

## ENDOMETRIAL CANCER AND miRNA

Endometrial cancer is the most frequent among malignant female genital cancers and approximately 80 % of cases have a histological type of endometrioid adenocarcinoma (Amant et al., 2005). The oncogenic pattern of endometrial cancer and factors for metastasis and invasion have been established and the expression pattern of miRNAs differs from that of normal endometrium (Table 1). Among endometrial cancers, endometrioid carcinoma and papillary carcinoma also have different miRNA expression (Chan et al., 2011).

In endometrial cancer, miRNAs including miR-185, miR-106a, miR-181a, miR-210, miR-423, miR-103, miR-107, miR-let7c, miR-205, miR-449 and miR-429 are upregulated and involved in oncogenesis, invasion and metastasis (Boren et al., 2008; Wu et al., 2009; Chung et al., 2009). miR-7 is also upregulated in endometrial cancer, and invasion and cancer cell migration are inhibited by downregulation of miR-7 using an anti-miRNA antibody (Chung et al., 2012). In endometrioid adenocarcinoma, up-

regulation of miR-27 expression is surgical stage-dependent. miR-27 contributes to survival of tumor cells through the resultant reduced expression of FOXO1, a target gene of miR-27 that inhibits apoptosis (Mozos et al., 2014).

Dai et al. (2013) found that miR-200b overexpression in adenocarcinoma cell lines inhibited expression of tissue inhibitor of metalloproteinase-2 (TIMP2) and increased the level of matrix metalloproteinase (MMP) 2, indicating that miR-200b is involved in metastasis of endometrial cancer. Six miRNAs show specific expression patterns in adenocarcinoma, including miR-34b, which is methylated in the promoter region in adenocarcinoma and is involved in proliferation and invasion (Hiroki et al., 2012). In contrast, multiple miRNAs including miR-let7e, miR-30c, miR-221, miR-152, miR-193, miR-204, miR-99b and miR-193b are downregulated in endometrial cancer (Boren et al., 2008; Wu et al., 2009; Chung et al., 2009). These miRNAs inhibit oncogenesis, invasion and metastasis, and these phenomena are induced by downregulation of the miRNAs.

**Table 1:** Changes in expression of miRNAs in endometrial cancer

Upregulation (OncomiRs)	Target genes	References
miR-185		Boren et al., 2008; Wu et al., 2009; Chung et al., 2009
miR-106a		
miR-181a		
miR-210		
miR-423		
miR-103		
miR-107		
miR-let7c		
miR-205		
miR-449		
miR-429		
miR-7		Chung et al., 2012
miR-27	<i>FOXO1</i>	Mozos et al., 2014
miR-200b	<i>TIMP2</i>	Dai et al., 2013
miR-34b		Hiroki et al., 2012
Downregulation (Tumor-suppressor miRs)	Target genes	References
miR-let7e		Boren et al., 2008; Wu et al., 2009; Chung et al., 2009
miR-221		
miR-193		
miR-99b		
miR-193b		
miR-204	<i>FOXO1</i>	Boren et al., 2008; Wu et al., 2009; Chung et al., 2009; Chung et al., 2012
miR-152	<i>DNMT1, E2F3, MET, Rictor</i>	Boren et al., 2008; Wu et al., 2009; Chung et al., 2009; Tsuruta et al., 2011; Banno et al., 2012
miR-30c	<i>MTA1</i>	Xie Y et al., 2011; Zhou H et al., 2012; Kong et al., 2014
miR-129-2	<i>SOX4, hMLH1</i>	Huang et al., 2011; Banno et al., 2012; Huang et al., 2014
miR-203		Huang et al., 2014
miR-106b	<i>TMIST1</i>	Dong et al., 2014
miR-let7a	<i>Aurora-B</i>	Liu et al., 2013

miRNAs are also involved in DNA methylation during progression of endometrial cancer. For example, if expression of miR-129-2 is enhanced by epigenetic mechanisms including DNA demethylation and histone acetylation, expression of SRY-related high-mobility group box 4 (SOX4) is suppressed and growth of endometrial cancer is inhibited. In endometrial cancer, this mechanism fails and SOX4 is overexpressed (Huang et al., 2009). miR-129-2 is involved in microsatellite instability and methylation of human mutL homolog 1 (hMLH1), a DNA mismatch repair (MMR) gene implicated in progression of type I endometrial cancer. hMLH1 is frequently methylated in endometrial cancer tissues and may induce mutation of cancer-associated genes including human mutS homolog 6 (hMSH6), type II transforming growth factor-beta (TGF- $\beta$ II), Bcl2-associated X protein (BAX) and PTEN (Banno et al., 2012). In addition to miR-129-2, miR-203 also regulates SOX4, and methylation of these miRNAs may lead to development of endometrial cancer (Huang et al., 2014).

Expression of miR-152 is regulated by methylation and is downregulated in endometrial cancer. miR-152 inhibits expression of targets including DNA methyltransferase (DNMT1), E2F3, met proto-oncogene (MET) and rapamycin-insensitive companion of mTOR (Rictor), all of which are implicated in oncogenesis. Thus, treatment targeting miR-152 and use of this miRNA as a biomarker have potential (Banno et al., 2012; Tsuruta et al., 2011). miR-106b inhibits EMT and TWIST1 (Dong et al., 2014) and metastasis is promoted by downregulation of miR-106b. let-7a inhibits Aurora-B and reduces the level of Aurora-B protein, with resultant inhibition of endometrial cancer onset (Liu et al., 2013). miR-204 also regulates metastasis and invasion of tumor cells in the endometrium. A target of miR-204, forkhead box C1 (FOXC1), regulates metastasis and invasion in human endometrial cancer-derived HEC1A cells. In endometrial cancer, FOXC1 is overexpressed due to downregulation of miR-204 and this promotes metastasis and

invasion of tumor cells (Chung et al., 2012).

miR-30c acts directly on metastasis-associated gene-1 (MTA1). miR-30c inhibits cell proliferation in the endometrium via regulation of MTA1, and thus downregulation of miR-30c may be involved in type I and II endometrial cancer (Xie et al., 2011; Zhou et al., 2012). miR-30c is also downregulated by estradiol (E2) in estrogen receptor (ER)-positive Ishikawa cells and ER-negative HEC1B cells, which indicates that estrogen regulates miR-30c in the endometrium and is involved in oncogenesis of endometrial cancer (Kong et al., 2014). Lin et al. (2008) showed that miR-302 induces demethylation of overall genomic DNA, and consequently activates transcription factors including Oct4, Sox2, Nanog and Lin28. Expression of specific genes in human ES cells causes reprogramming in these cells, with somatic cells changed to pluripotent stem (mirPS) cells. miR-302 inhibits tumorigenesis in various cancers, including through direct inhibition of cyclin D1 and indirect inhibition of CDK1 (Yan et al., 2014).

Many important findings have been described in this section, but it is also clear that an understanding of the mechanisms of miRNAs in endometrial cancer requires further studies.

#### **APPLICATION OF miRNA IN ENDOMETRIAL CANCER AS A BIOMARKER**

Valadi et al. reported the initial finding of extracellular miRNAs in exosomes (Valadi et al., 2007). Exosomes are extracellular vesicles and miRNA carriers that are released from cells into blood. Catabolic enzymes for miRNA are present in serum and it was originally thought that miRNAs could not exist in blood. However, miRNAs in exosomes are stable and are present in blood (Chim et al., 2008; Gilad et al., 2008). Exosomes released from cancer cells may have a relationship with mechanisms including immunosuppression, drug resistance and angiogenesis. Thus, miRNAs in exosomes reflect cell characteristics and overexpressed miRNAs in cancer cells are included in exosomes released from

these cells. Lawrie et al. (2008) first showed that cancer-specific miRNAs were effective biomarkers, and subsequently development of biomarkers using miRNAs has increased.

Several miRNAs have specific expression profiles in particular carcinomas and are likely to be useful as biomarkers. Analysis of miRNAs may classify cancer more correctly than analysis of 20,000 mRNAs (Lu et al., 2005). Lung cancer with downregulation of let-7 has a poor prognosis after surgery, indicating that miRNAs may be markers for prognosis, as well as for early diagnosis (Takamizawa et al., 2004). Most patients with endometrial cancer are diagnosed at an early clinical stage. However, the 5-year survival rate of patients in whom endometrial cancer is detected at an advanced stage is low, ranging from 10 % to 29 %. Therefore, development of a specific marker is required for early detection of endometrioid adenocarcinoma (Bansal et al., 2009). For example, Tan et al. (2010) found that upregulation of miR-155 was related to cancer stage and lymph node metastases in a study of miR-155 in serum that had the goal of detection of differences in miRNA expression among endometrioid adenocarcinoma, normal endometrium, and other tissue types.

Endometriosis is one of precancerous lesion of endometrial cancer that is a target for early diagnosis. A combination of miR-199a and miR-542-3p can be used to diagnose endometriosis with a sensitivity of 96.61 % and specificity of 79.66 %, and this approach may serve as a noninvasive marker in practice (Yu et al., 2012). Torres et al. (2012) found that expression of miR-99a, miR-100 and miR-199b was upregulated in serum of patients with endometrioid adenocarcinoma, and showed that a combination of miR-99a and miR-199b had a higher diagnostic value than each miRNA alone. Analysis of miRNAs in serum in a genome-wide study showed that a combination of four serum miRNAs, miR-222, miR-223, miR-186 and miR-204, can be used to diagnose endometrioid adenocarcinoma with high probability. Use of this combination for diagnosis gave an area under the ROC curve of 0.927, which

was higher than that for the currently used marker, CA-125 (0.673).

Type II endometrial cancer is difficult to diagnose because it involves de novo oncogenesis and has no precancerous lesion. However, several miRNAs including miR-125b may be useful for diagnosis of endometrial cancer. Expression of miR-125b in type II endometrial cancer cells is significantly upregulated compared to that in type I cells. Tumor protein p53 inducible nuclear protein 1 (TP53INP1) gene, a target of miR-125b, may be related to malignancy of type II endometrial cancer because cancer cells proliferate when this gene is not regulated (Jiang et al., 2011). V-erb-b2 erythroblastic leukemia viral oncogene homolog 2 (ERBB2), a second target of miR-125b, is also involved in invasion of cancer cells (Shang et al., 2012). miR-194 may also be a biomarker for prognostic diagnosis of endometrial cancer. miR-194 is markedly decreased in endometrial cancer and this decrease is correlated with the stage of cancer and is associated with prognosis (Zhai et al., 2013). Administration of pre-miRNA showed that miR-194 targets the oncogene BMI1 and inhibits the EMT phenotype and EC cell invasion (Dong et al., 2011).

KRAS mutation frequently occurs in type II endometrial cancer. Expression levels of miR-181b, miR-324-3p and miR-518b are downregulated in cancer with a KRAS mutation (Lee et al., 2014). Thus, miR-181b, miR-324-3p and miR-518b may be prognostic markers. miR-205 targeting PTEN is also associated with decreased survival. Expression profiles of miRNAs in endometrioid adenocarcinoma may also be implicated in clinicopathologic characteristics (Torres et al., 2013; Tsukamoto et al., 2014), with expression of several miRNAs showing a relationship with International Federation of Gynaecology and Obstetrics (FIGO) staging, cancer grade, recurrence and lymph node metastasis.

## APPLICATION OF miRNA FOR TREATMENT OF ENDOMETRIAL CANCER

miRNAs have a close relationship with cancer, and therefore may be useful for cancer treatment. Two approaches have been proposed as therapeutic strategies involving miRNAs. The first is replacement therapy with a particular miRNA for which downregulation is associated with disease onset. The second is replacement of molecules inhibiting a certain miRNA for which upregulation is associated with onset. The first approach includes treatment with viral vectors and ribosomes. miR-16 supplemented in bone metastasis model mice with prostate cancer reduced proliferation of local tumors and bone metastasis was significantly inhibited without adverse reactions. These results suggest that supplementation of downregulated miRNA can have a therapeutic effect (Takeshita et al., 2010). Similarly, administration of miR-143 inhibited metastasis in pulmonary metastasis model mice with osteosarcoma (Osaki et al., 2011). The second approach includes inhibition of a target miRNA using locked nucleic acid (LNA). Using this approach, activities of miR-122 in primates with hepatitis C were inhibited without serious adverse reactions and HCV proliferation was also inhibited (Randall et al., 2007; Lanford et al., 2010).

Studies of miRNAs have revealed detailed mechanisms of drugs. Such drugs include bortezomib, an inhibitor of ubiquitin-dependent proteolysis that kills tumor cells in vitro and in animal models. This drug blocks proteasome activity, arrests cells in G2/M phase, induces apoptosis, and inhibits development of endometrial cancer. The action of miR-17-5p on p21 increased the efficacy of bortezomib (Shen et al., 2013).

Drug sensitivity of cancer can also be influenced using miRNAs. In endometrial cancer, expression of miR-34c, which regulates metastasis, cell death and invasion, is markedly downregulated, and a combination of a miR-34c mimic with cisplatin improved the drug efficacy in cell lines (Jiang et al., 2013). Expression of miR-200b, miR-200c and miR-429, which inhibit transcription factor

AP-2 $\alpha$ , is upregulated in endometrial cancer and the rate of expression is positively correlated with resistance to cisplatin. However, SNP (rs1045385A>C) decreases the effect on AP-2 $\alpha$ , and sensitivity to cisplatin is increased (Wu et al., 2011). These studies suggest that a comprehensive investigation of which miRNA causes resistance or increases sensitivity to a particular drug will improve selection of drug treatment.

miRNAs themselves can be used for cancer treatment as a means of increasing expression of tumor suppressor genes and inhibiting oncogenes. Administration of a tumor suppressor miR, miR-152, in vitro and in vivo gave significant tumor suppression (Tsuruta et al., 2011). However, no miRNA is currently used in clinical practice, in part because application of miRNAs depends on effective and cancer cell-specific transport. miRNAs exist stably in various tissues and structures, and therefore should be relatively easy to transport into cells via blood vessels and other structures (Mitchell et al., 2008). Transport methods for short interfering RNA (siRNA) should also be applicable to miRNA (Gondi and Rao, 2009; Broderick and Zamore, 2011) due to the structural similarity of miRNA and siRNA. Nanoparticles contribute to the stability of miRNAs (Kong et al., 2012) and liposome-polycation-hyaluronic acid (LPH) modified with a single-chain antibody fragment (scFv) has been used for transport of miRNAs and siRNAs into cancer cells, with resultant downregulation of target genes (c-Myc, MDM2 and VEGF). Intravenous injection of nanoparticles modified with a humanized monoclonal antibody GC4 and miR-34a and two siRNAs also resulted in significant accumulation in cancer cells and inhibition of expression of target genes (Chen et al., 2010).

These results suggest that miRNAs can be transported specifically into cancer cells using nanoparticles modified with antibodies to cancer cell surface markers, and that a combination of siRNAs and miRNAs can increase the efficiency of gene regulation. Effects of miRNAs are relatively mild in comparison with molecular targeted drugs, but

miRNAs inhibit expression of multiple target molecules and cause fewer adverse reactions in normal cells.

### CONCLUSION

It is only 20 years since miRNAs were discovered, but many mechanisms have now been elucidated. Cancer cells include miRNAs with cancer-specific upregulation and downregulation, and these miRNAs are involved in oncogenesis, invasion and metastasis. miRNAs with expression patterns that differ from those in normal tissues, such as miR-125b, may be useful in treatment and as biomarkers for endometrial cancer. These miRNAs are currently being studied at an investigational level prior to development of clinical applications.

### Acknowledgments

We thank Dr. Y. Matsubara and Dr. E. Yuasa for helpful assistance. The authors gratefully acknowledge grant support from the Japan Society for the Promotion of Science (JSPS) through a Grant-in-Aid for Scientific Research (KAKENHI), a Grant-in-Aid for Scientific Research (C) (25462608), and a Grant-in-Aid for Young Scientists (B) (24791718); the Medical Research Encouragement Prize of The Japan Medical Association; and the Keio Gijyuku Academic Development Fund. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

### Conflict of Interest

The authors declare that they have no conflict of interest.

### REFERENCES

- Amant F, Moerman P, Neven P, Timmerman D, Van Limbergen E, Vergote I. Endometrial cancer. *Lancet*. 2005;366:491-505.
- Banno K, Kisu I, Yanokura M, Masuda K, Ueki A, Kobayashi Y, et al. Epigenetics and genetics in endometrial cancer: new carcinogenic mechanisms and relationship with clinical practice. *Epigenomics*. 2012; 4:147-62.
- Bansal N, Yendluri V, Wenham RM. The molecular biology of endometrial cancers and the implications for pathogenesis, classification, and targeted therapies. *Cancer Control*. 2009;16:8-13.
- Boren T, Xiong Y, Hakam A, Wenham R, Apte S, Wei Z, et al. MicroRNAs and their target messenger RNAs associated with endometrial carcinogenesis. *Gynecol Oncol*. 2008;110:206-15.
- Broderick JA, Zamore PD. MicroRNA therapeutics. *Gene Ther*. 2011;18:1104-10.
- Calin GA, Dumitru CD, Shimizu M, Bichi R, Zupo S, Noch E, et al. Frequent deletions and down-regulation of micro-RNA genes miR15 and miR16 at 13q14 in chronic lymphocytic leukemia. *Proc Natl Acad Sci U S A*. 2002;99:15524-9.
- Chan E, Prado DE, Weidhaas JB. Cancer microRNAs from subtype profiling to predictors of response to therapy. *Trends Mol Med*. 2011;17:235-43.
- Chen C, Ridzon D, Lee CT, Blake J, Sun Y, Strauss WM. Defining embryonic stem cell identity using differentiation-related microRNAs and their potential targets. *Mamm Genome*. 2007;18:316-27.
- Chen Y, Zhu X, Zhang X, Liu B, Huang L. Nanoparticles modified with tumor-targeting scFv deliver siRNA and miRNA for cancer therapy. *Mol Ther*. 2010; 18:1650-6.
- Chim SS, Shing TK, Hung EC, Leung TY, Lau TK, Chiu RW, et al. Detection and characterization of placental microRNAs in maternal plasma. *Clin Chem*. 2008;54:482-90.
- Chung TK, Cheung TH, Huen NY, Wong KW, Lo KW, Yim SF, et al. Dysregulated microRNAs and their predicted targets associated with endometrioid endometrial adenocarcinoma in Hong Kong women. *Int J Cancer*. 2009;124:1358-65.
- Chung TK, Lau TS, Cheung TH, Yim SF, Lo KW, Siu NS, et al. Dysregulation of microRNA-204 mediates migration and invasion of endometrial cancer by regulating FOXC1. *Int J Cancer*. 2012;130:1036-45.

- Dai Y, Xia W, Song T, Su X, Li J, Li S, et al. MicroRNA-200b is overexpressed in endometrial adenocarcinomas and enhances MMP2 activity by down-regulating TIMP2 in human endometrial cancer cell line HEC-1A cells. *Nucleic Acid Ther.* 2013;23:29-34.
- Dong P, Kaneuchi M, Watari H, Hamada J, Sudo S, Ju J, et al. MicroRNA-194 inhibits epithelial to mesenchymal transition of endometrial cancer cells by targeting oncogene BMI-1. *Mol Cancer.* 2011;10:99.
- Dong P, Kaneuchi M, Watari H, Sudo S, Sakuragi N. MicroRNA-106b modulates epithelial-mesenchymal transition by targeting TWIST1 in invasive endometrial cancer cell lines. *Mol Carcinog.* 2014;53:349-59.
- Elbashir SM, Harborth J, Lendeckel W, Yalcin A, Weber K, Tuschl T. Duplexes of 21-nucleotide RNAs mediate RNA interference in cultured mammalian cells. *Nature.* 2001;411:494-8.
- Fujita S, Ito T, Mizutani T, Minoguchi S, Yamamichi N, Sakurai K, et al. miR-21 Gene expression triggered by AP-1 is sustained through a double-negative feedback mechanism. *J Mol Biol.* 2008;378:492-504.
- Gilad S, Meiri E, Yogev Y, Benjamin S, Lebanony D, Yerushalmi N, et al. Serum microRNAs are promising novel biomarkers. *PLoS One.* 2008;3:e3148.
- Gondi CS, Rao JS. Concepts in *in vivo* siRNA delivery for cancer therapy. *J Cell Physiol.* 2009;220:285-91.
- Hayashita Y, Osada H, Tatematsu Y, Yamada H, Yanagisawa K, Tomida S, et al. A polycistronic microRNA cluster, miR-17-92, is overexpressed in human lung cancers and enhances cell proliferation. *Cancer Res.* 2005;65:9628-32.
- Hiroki E, Suzuki F, Akahira J, Nagase S, Ito K, Sugawara J, et al. MicroRNA-34b functions as a potential tumor suppressor in endometrial serous adenocarcinoma. *Int J Cancer.* 2012;131:E395-404.
- Huang YW, Liu JC, Deatherage DE, Luo J, Mutch DG, Goodfellow PJ, et al. Epigenetic repression of microRNA-129-2 leads to overexpression of SOX4 oncogene in endometrial cancer. *Cancer Res.* 2009;69:9038-46.
- Huang YW, Kuo CT, Chen JH, Goodfellow PJ, Huang TH, Rader JS, et al. Hypermethylation of miR-203 in endometrial carcinomas. *Gynecol Oncol.* 2014;133:340-5.
- International Human Genome Sequencing Consortium: Finishing the euchromatic sequence of the human genome. *Nature.* 2004;431:931-45.
- Jiang F, Liu T, He Y, Yan Q, Chen X, Wang H, et al. MiR-125b promotes proliferation and migration of type II endometrial carcinoma cells through targeting TP53INP1 tumor suppressor *in vitro* and *in vivo*. *BMC Cancer.* 2011;11:425.
- Jiang L, Meng W, Zeng J, Hu H, Lu L. MiR-34c oligonucleotide enhances chemosensitivity of Ishikawa cell to cisplatin by inducing apoptosis. *Cell Biol Int.* 2013;37:577-83.
- Johnson SM, Grosshans H, Shingara J, Byrom M, Jarvis R, Cheng A, et al. RAS is regulated by the let-7 microRNA family. *Cell.* 2005;120:635-47.
- Kong X, Xu X, Yan Y, Guo F, Li J, Hu Y, et al. Estrogen regulates the tumour suppressor MiRNA-30c and its target gene, MTA-1, in endometrial cancer. *PLoS One.* 2014;9:90810.
- Kong YW, Ferland-McCollough D, Jackson TJ, Bushnell M. microRNAs in cancer management. *Lancet Oncol.* 2012;13:249-58.
- Lanford RE, Hildebrandt-Eriksen ES, Petri A, Persson R, Lindow M, Munk ME, et al. Therapeutic silencing of microRNA-122 in primates with chronic hepatitis C virus infection. *Science.* 2010;327:198-201.
- Lawrie CH, Gal S, Dunlop HM, Pushkaran B, Liggins AP, Pulford K, et al. Detection of elevated levels of tumour-associated microRNAs in serum of patients with diffuse large B-cell lymphoma. *Br J Haematol.* 2008;141:672-5.
- Lee YS, Dutta A. The tumor suppressor microRNA let-7 represses the HMG2 oncogene. *Genes Dev.* 2007;21:1025-30.
- Lee LJ, Ratner E, Uduman M, Winter K, Boeke M, Greven KM, et al. The KRAS-variant and miRNA expression in RTOG endometrial cancer clinical trials 9708 and 9905. *PLoS One.* 2014;9:94167.
- Lin SL, Chang DC, Chang-Lin S, Lin CH, Wu DT, Chen DT, et al. Mir-302 reprograms human skin cancer cells into a pluripotent ES-cell-like state. *RNA.* 2008;14:2115-24.
- Liu P, Qi M, Ma C, Lao G, Liu Y, Liu Y, et al. Let7a inhibits the growth of endometrial carcinoma cells by targeting Aurora-B. *FEBS Lett.* 2013;587:2523-9.
- Lu J, Getz G, Miska EA, Alvarez-Saavedra E, Lamb J, Peck D, et al. MicroRNA expression profiles classify human cancers. *Nature.* 2005;435:834-8.
- Mitchell PS, Parkin RK, Kroh EM, Fritz BR, Wyman SK, Pogosova-Agadjanyan EL, et al. Circulating microRNAs as stable blood-based markers for cancer detection. *Proc Natl Acad Sci U S A* 2008;105:10513-8.

- Mozos A, Catasús L, D'Angelo E, Serrano E, Espinosa I, Ferrer I, et al. The FOXO1-miR27 tandem regulates myometrial invasion in endometrioid endometrial adenocarcinoma. *Hum Pathol.* 2014;45:942-51.
- Olive V, Bennett MJ, Walker JC, Ma C, Jiang I, Cordon-Cardo C, et al. miR-19 is a key oncogenic component of mir-17-92. *Genes Dev.* 2009;23:2839-49.
- Osaki M, Takeshita F, Sugimoto Y, Kosaka N, Yamamoto Y, Yoshioka Y, et al. MicroRNA-143 regulates human osteosarcoma metastasis by regulating matrix metalloproteinase-13 expression. *Mol Ther.* 2011;19:1123-30.
- Randall G, Panis M, Cooper JD, Tellinghuisen TL, Sukhodolets KE, Pfeffer S, et al. Cellular cofactors affecting hepatitis C virus infection and replication. *Proc Natl Acad Sci U S A* 2007;104:12884-9.
- Shang C, Lu YM, Meng LR. MicroRNA-125b down-regulation mediates endometrial cancer invasion by targeting ERBB2. *Med Sci Monit.* 2012;18:149-55.
- Shen Y, Lu L, Xu J, Meng W, Qing Y, Liu Y, et al. Bortezomib induces apoptosis of endometrial cancer cells through microRNA-17-5p by targeting p21. *Cell Biol Int.* 2013;37:1114-21.
- Stefani G, Slack FJ. Small non-coding RNAs in animal development. *Nat Rev Mol Cell Biol.* 2008;9:219-30.
- Takamizawa J, Konishi H, Yanagisawa K, Tomida S, Osada H, Endoh H, et al. Reduced expression of the let-7 microRNAs in human lung cancers in association with shortened postoperative survival. *Cancer Res.* 2004;64:3753-6.
- Takeshita F, Patrawala L, Osaki M, Takahashi RU, Yamamoto Y, Kosaka N, et al. Systemic delivery of synthetic microRNA-16 inhibits the growth of metastatic prostate tumors via downregulation of multiple cell-cycle genes. *Mol Ther.* 2010;18:181-7.
- Tan ZQ, Liu FX, Tang HL, Su Q. [Expression and its clinical significance of hsa-miR-155 in serum of endometrial cancer]. *Zhonghua Fu Chan Ke Za Zhi.* 2010;45:772-4. [in Chinese].
- Torres A, Torres K, Pesci A, Ceccarino M, Paszkowski T, Cassandrini P, et al. Deregulation of miR-100, miR-99a and miR-199b in tissues and plasma coexists with increased expression of mTor kinase in endometrioid endometrial carcinoma. *BMC Cancer.* 2012;12:369.
- Torres A, Torres K, Pesci A, Ceccaroni M, Paszkowski T, Cassandrini P, et al. Diagnostic and prognostic significance of miRNA signatures in tissues and plasma of endometrioid endometrial carcinoma patients. *Int J Cancer.* 2013;132:1633-45.
- Tsakamoto O, Miura K, Mishima H, Abe S, Kaneuchi M, Higashijima A, et al. Identification of endometrioid endometrial carcinoma-associated microRNAs in tissue and plasma. *Gynecol Oncol.* 2014;132:715-21.
- Tsuruta T, Kozaki K, Uesugi A, Furuta M, Hirasawa A, Imoto I, et al. miR-152 is a tumor suppressor microRNA that is silenced by DNA hypermethylation in endometrial cancer. *Cancer Res.* 2011;71:6450-62.
- Valadi H, Ekström K, Bossios A, Sjöstrand M, Lee JJ, Lötvall JO. Exosome-mediated transfer of mRNAs and microRNAs is a novel mechanism of genetic exchange between cells. *Nat Cell Biol.* 2007;9:654-9.
- Volinia S, Calin GA, Liu CG, Ambs S, Cimmino A, Petrocca F, et al. A microRNA expression signature of human solid tumors defines cancer gene targets. *Proc Natl Acad Sci U S A* 2006;103:2257-61.
- Wu W, Lin Z, Zhuang Z, Liang X. Expression profile of mammalian microRNAs in endometrioid adenocarcinoma. *Eur J Cancer Prev.* 2009;18:50-5.
- Wu Y, Xiao Y, Ding X, Zhuo Y, Ren P, Zhou C, et al. A miR-200b/200c/429-binding site polymorphism in the 3' untranslated region of the AP-2a gene is associated with cisplatin resistance. *PLoS One.* 2011;6:29043.
- Xie Y, Wang YL, Yu L, Hu Q, Ji L, Zhang Y, et al. Metformin promotes progesterone receptor expression via inhibition of mammalian target of rapamycin (mTOR) in endometrial cancer cells. *J Steroid Biochem Mol Biol.* 2011;126:113-20.
- Yan GJ, Yu F, Wang B, Zhou HJ, Ge QY, Su J, et al. MicroRNA miR-302 inhibits the tumorigenicity of endometrial cancer cells by suppression of Cyclin D1 and CDK1. *Cancer Lett.* 2014;345:39-47.
- Yu F, Yao H, Zhu P, Zhang X, Pan Q, Gong Y, et al. let-7 regulates self renewal and tumorigenicity of breast cancer cells. *Cell.* 2007;131:1109-23.
- Yu S, Liu Y, Wang J, Guo Z, Zhang Q, Yu F, et al. Circulating microRNA profiles as potential biomarkers for diagnosis of papillary thyroid carcinoma. *J Clin Endocrinol Metab.* 2012;97:2084-92.
- Zhai H, Karaayvaz M, Dong P, Sakuragi N, Ju J. Prognostic significance of miR-194 in endometrial cancer. *Biomark Res.* 2013;1:12.
- Zhou H, Xu X, Xun Q, Yu D, Ling J, Guo F, et al. miR-30c negatively regulates endometrial cancer cells by targeting metastasis-associated gene-1. *Oncol Rep.* 2012;27:807-12.