

Supplementary material to:

**EFFECTS OF TRICHOSTATIN A ON DRUG UPTAKE
TRANSPORTERS IN PRIMARY RAT HEPATOCYTE CULTURES**

Eva Ramboer*, Vera Rogiers, Tamara Vanhaecke[°], Mathieu Vinken[°]

In Vitro Toxicology and Dermato-cosmetology research group, Center for Pharmaceutical Research, Faculty of Medicine and Pharmacy, Vrije Universiteit Brussel


[°] equally contributing last authors

* Corresponding author: Eva Ramboer; In Vitro Toxicology and Dermato-cosmetology research group, Center for Pharmaceutical Research, Faculty of Medicine and Pharmacy, Vrije Universiteit Brussel, (VUB), Laarbeeklaan 103, B-1090 Brussels, Belgium.


E-mail: eramboer@vub.ac.be, +32.2.477.45.87

<http://dx.doi.org/10.17179/excli2015-220>

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>).



Supplementary Figure 1: Accumulation of radioactive substrates as a function of substrate concentration. Freshly isolated rat hepatocytes were cultivated in 6-well plates (9.6 cm²) as described in “2.2 Hepatocyte isolation and cultivation”. Day 4 non-treated hepatocyte cultures were incubated for 90 seconds with a range of substrate concentrations, namely 0,02-10 µM [3H] Taurocholic acid (Ntcp) and 0,0022-1 µM [3H] Estrone-3-sulfate (Oatp4) either at 37 °C (active and passive transport) or 4 °C (passive transport). Active transport, equivalent to substrate uptake values at 37 °C minus uptake values at 4 °C was determined (n=1, 3 technical repeats). ([3H], tritium; D, day; Ntcp, sodium taurocholate cotransporting polypeptide; Oatp4, organic anion transporting polypeptide 4)



Supplementary Figure 2: Accumulation of radioactive substrates as a function of incubation time. Freshly isolated rat hepatocytes were cultivated in 6-well plates (9.6 cm²) as described in “2.2 Hepatocyte isolation and cultivation”. Day 1 non-treated hepatocyte cultures were incubated for 0.5, 1, 2, 3, 5 and 7 minutes with 1 µCi/ml [3H] Taurocholic acid (0,2 µM) (Ntcp) or [3H] Estrone-3-sulfate (0,02 µM) (Oatp4) (n=1, 1 technical repeat) ([³H], tritium; D, day; Ntcp, sodium taurocholate cotransporting polypeptide; Oatp4, organic anion transporting polypeptide 4)