

Supplementary data to:

**IMMUNOREGULATORY, PROLIFERATIVE AND ANTI-OXIDANT
EFFECTS OF NANOCURCUMINOIDS ON ADIPOSE-DERIVED
MESENCHYMAL STEM CELLS**

Forouzan Yousefi^{a#}, Fahimeh Lavi Arab^{a#}, Mahmoud Reza Jaafari^{b,c}, Maryam Rastin^d,
Nafiseh Tabasi^d, Mahdi Hatamipour^e, Karim Nikkhah^f, Mahmoud Mahmoudi^{a,g*}


- ^a Immunology Research Center, Mashhad University of Medical Sciences, Mashhad, Iran
^b Biotechnology Research Center, Pharmaceutical Technology Institute, Mashhad University of Medical Sciences, Mashhad, Iran
^c Department of Pharmaceutical Nanotechnology, School of Pharmacy, Mashhad University of Medical Sciences, Mashhad, Iran
^d Immunology Research Center, BuAli Research Institute, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran
^e Nanotechnology Research Center, Mashhad University of Medical Sciences, Mashhad, Iran
^f Department of Neurology, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran
^g Department of Immunology, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran.

Co-first authors: F.Y. and F.L. contributed equally to this research.


* Corresponding author: Mahmoud Mahmoudi, Department of Immunology, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran. Tel: +98 9151156304, Fax: +98 5138022229, E-mail: MahmoudiM@mums.ac.ir

<http://dx.doi.org/10.17179/excli2019-1366>

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>).



Supplementary Figure 1: Flow cytometry analysis of superficial markers of AT-MSCs based on comparison with negative isotype control on 10,000 events. Histogram plots show that AT-MSCs are positive for expression of CD73, CD105 and CD90, but negative for CD14, CD34, CD31, CD45 and HLA-DR.



Supplementary Figure 2: Tri-lineage capacity of AT-MSCs for differentiation into adipocytes, osteocytes and chondrocytes established by specific-cell staining after 21 days in a specialized medium: (a) fibroblast-like shape of AT-MSCs before differentiation; (b) and after differentiation of AT-MSCs toward adipocytes with lipid vacuoles detected by Oil Red O staining; (c) mineralization potential of AT-MSCs with appearance of calcium deposits following alizarin red staining; (d) and after alkaline phosphatase staining; (e) GAGs and sulfated proteoglycans from chondroitin cryosections after staining with hematoxylin-eosin; (f) and after toluidine blue.