

Supplementary material to:

**[(7-CHLOROQUINOLIN-4-YL)AMINO]ACETOPHENONES AND
THEIR COPPER(II) DERIVATIVES:
SYNTHESIS, CHARACTERIZATION, COMPUTATIONAL STUDIES
AND ANTIMALARIAL ACTIVITY**

Yonathan de J. Parra^{a*}, Felix D. Andueza L^a, Rosa E. Ferrer M^b, Julia Bruno Colmenarez^c,
María E. Acosta^d, Jaime Charris^e, Philip J. Rosenthal^f, Jiri Gut^f

^a Escuela de Ingeniería Ambiental, Facultad de Ingeniería en Geología, Minas, Petróleos y Ambiental, Universidad Central del Ecuador, Quito 170521, Ecuador

^b Departamento de Química, Facultad de Humanidades y Educación, Universidad del Zulia, Apartado 526, Maracaibo, Estado Zulia, Venezuela

^c Centro de Investigación y Tecnología de Materiales (CITeMa), Instituto Venezolano de Investigaciones Científicas, Apartado 20632, Altos de Pipe 1020-A Estado Miranda, Venezuela

^d Laboratorio de Bioquímica, Facultad de Farmacia, Universidad Central de Venezuela, Apartado 47206, Los Chaguaramos, 1041-A Caracas, Venezuela


^e Laboratorio de Síntesis Orgánica, Facultad de Farmacia, Universidad Central de Venezuela, Apartado 47206, Los Chaguaramos, 1041-A Caracas, Venezuela

^f Department of Medicine, University of California, Box 0811, San Francisco, California 94143, USA


* **Corresponding author:** Prof. Yonathan de J. Parra, Escuela de Ingeniería Ambiental, Facultad de Ingeniería en Geología, Minas, Petróleos y Ambiental, Universidad Central del Ecuador, Quito 170521, Ecuador; Tel: +593-2-2550588, ext. 111; E-mail: ydparra@uce.edu.ec

<http://dx.doi.org/10.17179/excli2019-1805>


This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>).




Supplementary Figure 1: ^1H NMR spectrum of 4-[(7-chloroquinolin-4-yl)amino]acetophenone (**4**)




Supplementary Figure 2: ^1H NMR spectrum of {4-[(7-chloroquinolin-4-yl)amino]acetophenone} copper (II) chloride (**4a**)




Supplementary Figure 3: Amplified ^1H NMR spectrum of 4-[(7-chloroquinolin-4-yl)amino]acetophenone (**4**)




Supplementary Figure 4: Amplified ^1H NMR spectrum of {4-[(7-chloroquinolin-4-yl)amino]acetophenone}copper(II) chloride (**4a**)




Supplementary Figure 5: ^{13}C NMR spectrum of 4-[(7-chloroquinolin-4-yl)amino]acetophenone (4)




Supplementary Figure 6: ^{13}C NMR spectrum of {4-[(7-chloroquinolin-4-yl)amino]acetophenone}copper(II) chloride (4a)




Supplementary Figure 7: ¹H NMR spectrum of 3-[(7-chloroquinolin-4-yl)amino]acetophenone (5)




Supplementary Figure 8: ¹H NMR spectrum of {3-[(7-chloroquinolin-4-yl)amino]acetophenone} copper(II) chloride (5a)




Supplementary Figure 9: Amplified ^1H NMR spectrum of 3-[(7-chloroquinolin-4-yl)amino]acetophenone (5)




Supplementary Figure 10: Amplified ^1H NMR spectrum of {3-[(7-chloroquinolin-4-yl)amino]acetophenone}copper(II) chloride (5a)




Supplementary Figure 11: ^{13}C NMR spectrum of 3-[(7-chloroquinolin-4-yl)amino]acetophenone (5)




Supplementary Figure 12: ^{13}C NMR spectrum of {3-[(7-chloroquinolin-4-yl)amino]acetophenone}copper(II) chloride (5a)




Supplementary Figure 13: Dissociative heterolytic mechanism for the a1, b1, b2, c1, d1 pathways; from the fragment at m/z 253-254 generated by the precursor molecular ion at m/z 297 for compound **4**. (Analogue mechanism for compound **5**)




Supplementary Figure 14: Dissociative homolytic mechanism for a2 pathway




Supplementary Figure 15: Dissociative heterolytic mechanism for the a4 and b3 pathways




Supplementary Figure 16: Dissociative heterolytic mechanism for the α_5 pathway



Supplementary Figure 17: Minor fragmentation channel of the cationic fragment $[4a]^+$ at m/z 657



Supplementary Figure 18: Extensive dissociation of the cationic fragment $[4a]^+$ at m/z 657



Supplementary Figure 19: Homolytic dissociation and reversible re-coordination/de-coordination of a water molecule for the fragment at m/z 359-361, derived from the ion $[4a]^+$ at m/z 657