

## Letter to the editor:

### RECENT UPDATES ON DAIDZEIN AGAINST OXIDATIVE STRESS AND CANCER

Sarita Rawat, Sachchidanand Pathak, Gaurav Gupta\*, Santosh Kumar Singh, Himmat Singh, Anurag Mishra, Ritu Gilhotra

School of Pharmacy, Suresh Gyan Vihar University, Mahal Road, Jagatpura, Jaipur, India

\* **Corresponding author:** Gaurav Gupta, School of Pharmacy, Suresh Gyan Vihar University, Mahal Road, Jagatpura, Jaipur, India. E-mail: [gauravpharma25@gmail.com](mailto:gauravpharma25@gmail.com)

<http://dx.doi.org/10.17179/excli2019-1847>

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>).

#### *Dear Editor,*

Daidzein (7-hydroxy-3-(4-hydroxyphenyl)-4H-chromen-4-one) is a naturally occurring compound commonly found in soybeans and some other legumes. Daidzein is an isoflavone by nature and isolated from *Pueraria Mirifica*, having category of biologically active secondary metabolites commonly produced in the soybean growth and belong to the group of flavonoids. A number of pharmacological activities have been accounted for daidzein, which includes anti-carcinogenesis, anti-fibrotic, anti-diabetic, cholesterol-lowering and cardiovascular activity. Daidzein pretreatment was found to diminish the seriousness of mucosal damage in a portion subordinate way. Some other therapeutic uses of daidzein are weight reduction, decreasing bowels moment, and inflammation associated with histopathological deformities (Amaral et al., 2017). Their structure and function are alike to human estrogen, which can play a massive role in the prevention of osteoporosis, cancer, and postmenopausal syndromes. Uses of daidzein in anticancer activity against ovarian cancer are still limited (Meng et al., 2017). Epidemiological data suggest that increased utilization of soybean in food results in decreased cancer risk. Soybean food is highly recommended in cancer prevention because it has a number of anticarcinogens (Yu et al., 2017). The potent antioxidant activity of daidzein reported in various *in vitro* and *in vivo* studies was conclusively shown in this literature (Table 1).

**Table 1:** Recent updates on daidzein against oxidative stress and cancer

Key findings	Reference
Daidzein could inhibit 5-FU-induced intestinal mucositis through prevention of relevant clinical symptoms, stimulation of the proliferative activity in the epithelium and crypts, suppression of oxidative stress and regulation of inflammatory homeostasis.	Atiq et al., 2019
Soluble CXCL16 and VEGF-A are two promoters of angiogenesis and metastasis in breast cancer. The stimulation of these two angiogenesis-related cytokines might represent one of the mechanisms explaining the proliferative effects of low isoflavone doses in estrogen-dependent cells. In estrogen-independent cells, daidzein inhibited their secretion, indicating anti-angiogenic properties.	Uifalean et al., 2018
The utilization of daidzein is associated with a reduced risk of prostate cancer but the direct effects of daidzein on the androgen receptor (AR) signaling pathway are not well understood.	Sivoňová et al., 2019
Cytotoxic effects of daidzein on both MIA PaCa-2 and HT-29 cell lines occurred in a dose-dependent manner. Daidzein exhibits genotoxic potential under <i>in vitro</i> conditions.	Gundogdu et al., 2018
Cell models and clinically possible concentrations of soy isoflavones do not support the notion that daidzein or other soy isoflavones can increase the effects of taxane chemotherapy in CRPC cell and xenograft models.	Eskra et al., 2019
Genistein and daidzein inhibit the accumulation of lipid droplets (LDs) by regulating LDs-related factors and lead to final apoptosis of colon cancer cells and possible molecular mechanisms of isoflavones in anti-obesity and anti-tumor functions.	Liang et al., 2018
Genistein, daidzein, and ERB-041 decreased ovarian cancer cell migration, invasion, proliferation, and bubble formation, signifying their roles on ovarian cancer cell metastasis, tumorigenesis, and their potential as other therapies for ovarian cancer patients.	Chan et al., 2018
Daidzein exerted neuroprotective effect against ICV-STZ-induced behavioral parameters include cognitive deficits and locomotor impairment. Where daidzein was able to correct oxidative stress in neuronal cells in the brain.	Wei et al., 2019
Daidzein enhanced cisplatin-impaired antioxidant defense such as reduced glutathione reserve, GPX activity, and SOD activity.	Meng et al., 2017
Daidzein attenuates CP-induced oxidative stress on blood cells and antioxidants in rats.	Karale and Kamath, 2017
Daidzein reduces hyperglycemia-induced oxidative stress in HUVECs. Thus, daidzein may be a potential therapeutic agent that could reduce hyperglycemia-induced oxidative damage associated with diabetes.	Park et al., 2016
Daidzein dosing reduced the B cell population as well as IgG1 production, while increased the helper and cytotoxic T cell population in females. The shortage of protection or exacerbation of T1D can be explained by the reduced B cell population and increased regulatory cells, revealing a completely unusual mechanism compared to GEN exposure.	Huang et al., 2019

**Table 1 (cont.):** Recent updates on daidzein against oxidative stress and cancer

Key findings	Reference
Considering the activity of daidzein as nutraceutical that forms complexes with actin that are helpful in the prevention of cancer metastases.	Budryn et al., 2018
Anti-proliferative action of 12 anticarcinogens from soy-bean against MCF-7 and MDA-MB-231 human breast cancer cell outline by single or two-way combination treatment. There is a synergistic result of the combination treatment of genistein plus daidzein MCF-7 cells.	Zhu et al., 2018
Daidzein exerts similar activity against SKOV3 cells in nude mouse xenograft models and it was shown that daidzein considerably reduced the tumorigenesis in vivo, indicative of the potential for daidzein as a lead molecule in the growth of ovarian cancer chemotherapy.	Hua et al., 2018
Daidzein may induce apoptosis of choriocarcinoma cells in a dose-dependent manner via the mitochondrial apoptotic pathway.	Zheng et al., 2018
Daidzein and equol do not provoke a major up-regulation of the transporter expression but instead an inhibition of BCRP activity and sensitization to BCRP substrates.	Rigalli et al., 2019
E1 and E2 conjugation by low micromolar concentrations of daidzein in the human breast cancer cell line MCF7, which leads to a small but statistically significant increase in unconjugated E2 of approximately 20 %.	Poschner et al., 2017
The impact of the dietary isoflavone daidzein on the biochemical fingerprint of two human breast cancer cell lines – estrogen-responsive (MCF-7) and estrogen-unresponsive- was found to induce a noticeable decrease in cell growth.	Medeiros et al., 2016
Daidzein is just as useful as genistein in protecting cells against oxidative damage especially with respect to DNA. Antioxidant activity of isoflavones could really contribute to the healthy properties of soy.	Foti et al., 2005
Oxidative DNA damage by daidzein metabolites plays a role in tumor initiation and that cell proliferation by isoflavones via ER-ERE binding induces tumor support and/or progression, resulting in cancer of estrogen-sensitive organs.	Murata et al., 2004
Antioxidant enzyme (AOE) system provoked by daidzein affected the oxidant rather than the antioxidant properties of daidzein.	Rohrdanz et al., 2002
Daidzein supplementation may protect cells from oxidative stress-inducing agents by inhibiting NF-kappa B activation and decreasing DNA adduct levels.	Davis et al., 2001
Daidzein can inhibit choriocarcinoma cell proliferation <i>in vitro</i> and <i>in vivo</i> ; the underlying mechanism behind the inhibitory effects may probably be suppressing ERK pathway and afterward arresting cell cycle at G1 phase.	Zheng et al., 2017
Biotransformation increased the anti-aromatase activity and the anti-tumoral value of daidzein in breast cancer cells. Moreover, it clarifies the potential use of daidzein in the prevention and/or treatment of ER+ breast cancer.	Amaral et al., 2017

**Conflict of interest**

The authors declare no conflict of interest.

**REFERENCES**

- Amaral C, Toloi MRT, Vasconcelos LD, Fonseca MJV, Correia-da-Silva G, Teixeira N. The role of soybean extracts and isoflavones in hormone-dependent breast cancer: Aromatase activity and biological effects. *Food Funct.* 2017;8:3064-74.
- Atiq A, Shal B, Naveed M, Khan A, Ali J, Zeeshan S, et al. Daidzein ameliorates 5-fluorouracil-induced intestinal mucositis by suppressing oxidative stress and inflammatory mediators in rodents. *Eur J Pharmacol.* 2019;843:292-306.
- Budryn G, Grzelczyk J, Pérez-Sánchez H. Binding of red clover isoflavones to actin as a potential mechanism of anti-metastatic activity restricting the migration of cancer cells. *Molecules.* 2018;23:E2471.
- Chan KKL, Siu MKY, Jiang YX, Wang JJ, Leung THY, Ngan HYS. Estrogen receptor modulators genistein, daidzein and erb-041 inhibit cell migration, invasion, proliferation and sphere formation via modulation of fak and pi3k/akt signaling in ovarian cancer. *Cancer Cell Int.* 2018;18:65.
- Davis JN, Kucuk O, Djuric Z, Sarkar FH. Soy isoflavone supplementation in healthy men prevents nf-kappa b activation by tnf-alpha in blood lymphocytes. *Free Radic Biol Med.* 2001;30:1293-302.
- Eskra JN, Schlicht MJ, Bosland MC. Lack of combination effects of soy isoflavones and taxane chemotherapy of castration-resistant prostate cancer. *The Prostate.* 2019;79:223-33.
- Foti P, Erba D, Riso P, Spadafranca A, Criscuoli F, Testolin G. Comparison between daidzein and genistein antioxidant activity in primary and cancer lymphocytes. *Arch Biochem Biophys.* 2005;433:421-7.
- Gundogdu G, Dodurga Y, Cetin M, Secme M, Cicek B. The cytotoxic and genotoxic effects of daidzein on mia paca-2 human pancreatic carcinoma cells and ht-29 human colon cancer cells. *Drug Chem Toxicol.* 2018;1-7;epub ahead of print.
- Hua F, Li CH, Chen XG, Liu XP. Daidzein exerts anticancer activity towards skov3 human ovarian cancer cells by inducing apoptosis and cell cycle arrest, and inhibiting the raf/mek/erk cascade. *Int J Mol Med.* 2018;41:3485-92.
- Huang G, Xu J, Guo TL. Isoflavone daidzein regulates immune responses in the b6c3f1 and non-obese diabetic (nod) mice. *Int Immunopharmacol.* 2019;71:277-84.
- Karale S, Kamath JV. Effect of daidzein on cisplatin-induced hematotoxicity and hepatotoxicity in experimental rats. *Indian J Pharmacol.* 2017;49:49-54.
- Liang YS, Qi WT, Guo W, Wang CL, Hu ZB, Li AK. Genistein and daidzein induce apoptosis of colon cancer cells by inhibiting the accumulation of lipid droplets. *Food Nutr Res.* 2018;62:10.29219/fnr.v62.1384.
- Medeiros PS, Batista de Carvalho AL, Ruano C, Otero JC, Marques MP. Raman microspectroscopy for probing the impact of a dietary antioxidant on human breast cancer cells. *Food Funct.* 2016;7:2800-10.
- Meng H, Fu G, Shen J, Shen K, Xu Z, Wang Y, et al. Ameliorative effect of daidzein on cisplatin-induced nephrotoxicity in mice via modulation of inflammation, oxidative stress, and cell death. *Oxid Med Cell Longev.* 2017;2017:3140680.
- Murata M, Midorikawa K, Koh M, Umezawa K, Kawanishi S. Genistein and daidzein induce cell proliferation and their metabolites cause oxidative DNA damage in relation to isoflavone-induced cancer of estrogen-sensitive organs. *Biochemistry.* 2004;43:2569-77.
- Park MH, Ju JW, Kim M, Han JS. The protective effect of daidzein on high glucose-induced oxidative stress in human umbilical vein endothelial cells. *Z Naturforsch. C.* 2016;71(1-2):21-8.
- Poschner S, Maier-Salamon A, Zehl M, Wackerlig J, Dobusch D, Pachmann B, et al. The impacts of genistein and daidzein on estrogen conjugations in human breast cancer cells: A targeted metabolomics approach. *Front Pharmacol.* 2017;8:699.
- Rigalli JP, Scholz PN, Tocchetti GN, Ruiz ML, Weiss J. The phytoestrogens daidzein and equol inhibit the drug transporter bcrp/abcg2 in breast cancer cells: Potential chemosensitizing effect. *Eur J Nutr.* 2019;58:139-50.
- Rohrdanz E, Ohler S, Tran-Thi QH, Kahl R. The phytoestrogen daidzein affects the antioxidant enzyme system of rat hepatoma h4iie cells. *J Nutr.* 2002;132:370-5.
- Sivoňová MK, Kaplán P, Tatarková Z, Lichardusová L, Dušenka R, Jurečeková J. Androgen receptor and soy isoflavones in prostate cancer. *Mol Clin Oncol.* 2019;10:191-204.

Uifalean A, Rath H, Hammer E, Ionescu C, Iuga CA, Lalk M. Influence of soy isoflavones in breast cancer angiogenesis: A multiplex glass elisa approach. *J BUON*. 2018;23(7):53-9.

Wei J, Yang F, Gong C, Shi X, Wang G. Protective effect of daidzein against streptozotocin-induced alzheimer's disease via improving cognitive dysfunction and oxidative stress in rat model. *J Biochem Mol Toxicol*. 2019;33:e22319.

Yu B, Tang DZ, Li SY, Wu Y, Chen M. Daidzein promotes proliferation and differentiation in osteoblastic oct1 cells via activation of the bmp-2/smads pathway. *Pharmazie*. 2017;72(1):35-40.

Zheng W, Sun R, Yang L, Zeng X, Xue Y, An R. Daidzein inhibits choriocarcinoma proliferation by arresting cell cycle at g1 phase through suppressing erk pathway in vitro and in vivo. *Oncol Rep*. 2017;38:2518-24.

Zheng W, Liu T, Sun R, Yang L, An R, Xue Y. Daidzein induces choriocarcinoma cell apoptosis in a dose-dependent manner via the mitochondrial apoptotic pathway. *Mol Med Rep*. 2018;17:6093-9.

Zhu Y, Yao Y, Shi Z, Everaert N, Ren G. Synergistic effect of bioactive anticarcinogens from soybean on anti-proliferative activity in mda-mb-231 and mcf-7 human breast cancer cells in vitro. *Molecules*. 2018;23(7): E1557.