

Supplementary information to:

***POLYGONUM MINUS* ESSENTIAL OIL MODULATES CISPLATIN-INDUCED HEPATOTOXICITY THROUGH INFLAMMATORY AND APOPTOTIC PATHWAYS**

Norhashima Abd Rashid^{1¶}, Farida Hussan^{2¶}, Asmah Hamid^{3¶}, Nurul Raudzah Adib Ridzuan^{4¶}, Syarifah Aisyah Syed Abd Halim^{5¶}, Nahdia Afifah Abdul Jalil^{5¶}, Nor Haliza Mohamad Najib^{5¶}, Seong Lin Teoh^{5¶}, Siti Balkis Budin^{1*}


- ¹ Center for Diagnostic, Therapeutic and Investigative Studies, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, Kuala Lumpur, Malaysia
- ² Human Biology Department, School of Medicine, International Medical University, Bukit Jalil, Kuala Lumpur, Malaysia
- ³ Centre for Toxicology and Health Risk Studies, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, Kuala Lumpur, Malaysia
- ⁴ Department of Anatomy, Faculty of Medicine, Universiti Teknologi MARA, Selangor, Malaysia
- ⁵ Department of Anatomy, Faculty of Medicine, Universiti Kebangsaan Malaysia Medical Centre, Kuala Lumpur, Malaysia

¶ These authors contributed equally to this work.


* **Corresponding author:** Siti Balkis Budin, Center for Diagnostic, Therapeutic and Investigative Studies, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, Jalan Raja Muda Abdul Aziz, 50300 Kuala Lumpur, Malaysia. E-mail: balkis@ukm.edu.my

<http://dx.doi.org/10.17179/excli2020-2355>

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>).



Supplementary Figure 1: Flowchart of the effect of *P. minus* essential oil against cisplatin-induced hepatotoxicity



Supplementary Figure 2: MS chromatogram of *P. minus* essential oil

Supplementary Table 1: The mean and SEM of the effect of *P. minus* essential oil on CYP2E1 gene expression level in the liver of cisplatin-induced hepatotoxicity rats

Groups	Relative mRNA expression
	Mean \pm SEM
Control (Normal Saline)	1.00 \pm 0.10
Cisplatin (10 mg/kg)	1.88 \pm 0.24 ^a
BCP (150 mg/kg)	1.04 \pm 0.11 ^b
PmEO100CP (<i>P. minus</i> Essential Oil 100 mg/kg + Cisplatin 10 mg/kg)	1.14 \pm 0.07 ^b
PmEO200CP (<i>P. minus</i> Essential Oil 200 mg/kg + Cisplatin 10 mg/kg)	1.43 \pm 0.10 ^b
PmEO400CP (<i>P. minus</i> Essential Oil 400 mg/kg + Cisplatin 10 mg/kg)	2.40 \pm 0.34 ^{abcde}
PmEO400 (<i>P. minus</i> Essential Oil 400 mg/kg)	1.74 \pm 0.06 ^{acdf}

Values are mean \pm SEM (n = 6).

^a Significantly different from control group at p<0.05.

^b Significantly different from cisplatin group p<0.05.

^c Significantly different from BCP group at p<0.05.

^d Significantly different from PmEO100CP group at p<0.05.

^e Significantly different from PmEO200CP group at p<0.05.

^f Significantly different from PmEO400CP group at p<0.05.

Supplementary Table 2: The mean and SEM of the effect of *P. minus* essential oil on inflammation markers in the liver of cisplatin-induced hepatotoxicity rats

Groups	TNF- α	IL-1 α	IL-1 β	IL-6	IL-10
	Mean \pm SEM	Mean \pm SEM	Mean \pm SEM	Mean \pm SEM	Mean \pm SEM
Control (Normal Saline)	3.14 \pm 0.30	138.01 \pm 4.6	563.17 \pm 35.35	3661.51 \pm 208.30	912.31 \pm 65.04
Cisplatin (10 mg/kg)	7.20 \pm 0.37 ^a	357.22 \pm 10.91 ^a	915.30 \pm 32.87 ^a	5030.50 \pm 199.94 ^a	636.46 \pm 18.20 ^a
BCP (150 mg/kg)	4.47 \pm 0.54 ^b	151.82 \pm 6.93 ^b	676.74 \pm 15.40 ^b	3854.20 \pm 136.29 ^b	972.98 \pm 59.44 ^b
PmEO100CP (<i>P. minus</i> Essential Oil 100 mg/kg + Cisplatin 10 mg/kg)	4.01 \pm 0.70 ^b	194.10 \pm 24.83 ^b	591.46 \pm 74.60 ^b	3390.46 \pm 408.90 ^b	878.60 \pm 68.90 ^b
PmEO200CP (<i>P. minus</i> Essential Oil 200 mg/kg + Cisplatin 10 mg/kg)	6.30 \pm 0.37 ^a	298.24 \pm 22.45 ^{acd}	865.25 \pm 60.06 ^{ad}	6001.87 \pm 197.33 ^{acd}	576.75 \pm 54.76 ^{acd}
PmEO400CP (<i>P. minus</i> Essential Oil 400 mg/kg + Cisplatin 10 mg/kg)	9.42 \pm 1.08 ^{acde}	304.98 \pm 26.91 ^{acd}	828.90 \pm 56.13 ^{ad}	4849.26 \pm 108.71 ^{ade}	500.82 \pm 42.48 ^{acd}
PmEO400 (<i>P. minus</i> Essential Oil 400 mg/kg)	3.35 \pm 0.48 ^{bef}	191.96 \pm 11.81 ^{bef}	553.41 \pm 48.05 ^{bef}	3431.50 \pm 291.75 ^{bef}	504.68 \pm 34.27 ^{acd}

Values are mean \pm SEM (n=6).

^a Significantly different from control group at p<0.05.

^b Significantly different from cisplatin group at p<0.05.

^c Significantly different from BCP group at p<0.05.

^d Significantly different from PmEO100CP group at p<0.05.

^e Significantly different from PmEO200CP group at p<0.05.

^f Significantly different from PmEO400CP group at p<0.05.

Supplementary Table 3: The mean and SEM of the effect of *P. minus* essential oil on apoptosis gene expression in the liver of cisplatin-induced hepatotoxicity rats

Groups	Relative mRNA expression						
	p53	Caspase 8	Bax	Apaf-1	Caspase 9	Caspase 3	Bcl-2
	Mean ± SEM	Mean ± SEM	Mean ± SEM	Mean ± SEM	Mean ± SEM	Mean ± SEM	Mean ± SEM
Control (Normal Saline)	1.00 ± 0.02	1.00 ± 0.00	1.00 ± 0.00	1.00 ± 0.00	1.00 ± 0.00	1.00 ± 0.00	1.00 ± 0.10
Cisplatin (10 mg/kg)	2.00 ± 0.02 ^a	2.51 ± 0.00 ^a	1.67 ± 0.00 ^a	2.80 ± 0.00 ^a	4.94 ± 0.00 ^a	6.26 ± 0.06 ^a	0.25 ± 0.00 ^a
BCP (150 mg/kg)	0.64 ± 0.01 ^b	1.19 ± 0.01 ^b	1.31 ± 0.00 ^b	1.31 ± 0.00 ^b	1.11 ± 0.00 ^b	2.98 ± 0.09 ^b	0.89 ± 0.00 ^b
PmEO100CP (<i>P. minus</i> Essential Oil 100 mg/kg + Cisplatin 10 mg/kg)	0.59 ± 0.01 ^b	1.36 ± 0.00 ^b	1.09 ± 0.00 ^b	1.32 ± 0.00 ^b	1.22 ± 0.00 ^b	2.50 ± 0.05 ^b	0.85 ± 0.00 ^b
PmEO200CP (<i>P. minus</i> Essential Oil 200 mg/kg + Cisplatin 10 mg/kg)	0.82 ± 0.01 ^b	1.84 ± 0.02	1.86 ± 0.00 ^{acd}	2.00 ± 0.01	2.27 ± 0.00	5.25 ± 0.04 ^{ad}	0.34 ± 0.00 ^a
PmEO400CP (<i>P. minus</i> Essential Oil 400 mg/kg + Cisplatin 10 mg/kg)	2.96 ± 0.04 ^{acde}	2.42 ± 0.02 ^{ac}	1.87 ± 0.00 ^{acd}	4.56 ± 0.04 ^{abcde}	6.07 ± 0.02 ^{acde}	7.99 ± 0.07 ^{acde}	0.05 ± 0.00 ^{acd}
PmEO400 (<i>P. minus</i> Essential Oil 400 mg/kg)	0.80 ± 0.07 ^{bf}	1.60 ± 0.01	1.22 ± 0.00 ^{ef}	2.55 ± 0.01 ^{bef}	0.96 ± 0.00 ^{bf}	1.01 ± 0.03 ^{bef}	0.39 ± 0.00 ^a

Values are mean ± SEM (n = 6).

^a Significantly different from control group at p<0.05.

^b Significantly different from cisplatin group at p<0.05.

^c Significantly different from BCP group at p<0.05.

^d Significantly different from PmEO100CP group at p<0.05.

^e Significantly different from PmEO200CP group at p<0.05.

^f Significantly different from PmEO400CP group at p<0.05.

Supplementary Table 4: The mean and SEM of the effect of *P. minus* essential oil on cytochrome c protein expression in the liver of cisplatin-induced hepatotoxicity rats

Groups	Expression Level (Fold Change)
	Mean \pm SEM
Control (Normal Saline)	1.00 \pm 0.05
Cisplatin (10 mg/kg)	2.18 \pm 0.01 ^a
BCP (150 mg/kg)	1.18 \pm 0.00 ^b
PmEO100CP (<i>P. minus</i> Essential Oil 100 mg/kg + Cisplatin 10 mg/kg)	0.95 \pm 0.02 ^b
PmEO200CP (<i>P. minus</i> Essential Oil 200 mg/kg + Cisplatin 10 mg/kg)	1.86 \pm 0.02 ^{ad}
PmEO400CP (<i>P. minus</i> Essential Oil 400 mg/kg + Cisplatin 10 mg/kg)	2.31 \pm 0.03 ^{acd}
PmEO400 (<i>P. minus</i> Essential Oil 400 mg/kg)	1.68 \pm 0.05 ^{ad}

^aSignificantly different from control group at p<0.05.

^bSignificantly different from cisplatin group at p<0.05.

^cSignificantly different from BCP group at p<0.05.

^dSignificantly different from PmEO100CP group at p<0.05.