

Supplementary information to:

ROLE OF CHANGES IN SARS-COV-2 SPIKE PROTEIN IN THE INTERACTION WITH THE HUMAN ACE2 RECEPTOR: AN *IN SILICO* ANALYSIS

Joseph Thomas Ortega¹, Maria Luisa Serrano², Flor Helene Pujol³, Hector Rafael Rangel^{3*}

1 Department of Pharmacology and Cleveland Center for Membrane and Structural Biology, School of Medicine, Case Western Reserve University, Cleveland, OH 44106, USA


2 Unidad de Química Medicinal, Facultad de Farmacia, Universidad Central de Venezuela, Caracas, Venezuela

3 Laboratorio de Virología Molecular, Centro de Microbiología y Biología Celular, Instituto Venezolano de Investigaciones Científicas, Caracas, Venezuela

* **Corresponding author:** Hector Rafael Rangel, Laboratorio de Virología Molecular, Centro de Microbiología y Biología Celular, Instituto Venezolano de Investigaciones Científicas, Caracas, Venezuela; E-mail: hrangel2006@gmail.com

<http://dx.doi.org/10.17179/excli2020-1167>

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>).



Supplementary Figure 1: Spike protein of *Rhinolophus sinicus* (A) and the merge between spike of *Rhinolophus sinicus* and SARS-CoV (B). The merge image shows a high level of structural similarity in both protein, suggesting a similar pattern of interaction with the receptor.