

Original article:

**MICROBIOLOGICAL AND VIRULENCE ASPECTS OF
*RHODOTORULA MUCILAGINOSA***

Isabele Carrilho Jarros¹, Flávia Franco Veiga¹, Jakeline Luiz Corrêa¹, Isabella Letícia Esteves Barros¹, Marina Cristina Gadelha¹, Morgana F. Voidaleski², Neli Pieralisi³, Raissa Bocchi Pedroso¹, Vânia A. Vicente², Melyssa Negri¹, Terezinha Inez Estivalet Svidzinski^{1*}

¹ Division of Medical Mycology, Teaching and Research Laboratory in Clinical Analyses – Department of Clinical Analysis of State University of Maringá, Paraná, Brazil

² Postgraduate Program in Microbiology, Parasitology, and Pathology, Biological Sciences, Department of Basic Pathology, Federal University of Parana, Curitiba, Brazil

³ Department of Dentistry, State University of Maringá, Maringá, Paraná, Brazil

* **Corresponding author:** Terezinha Inez Estivalet Svidzinski, Division of Medical Mycology, Teaching and Research Laboratory in Clinical Analysis – Department of Clinical Analysis of State University of Maringá, Paraná, Brazil. Av. Colombo, 5790 CEP: 87020-900. Maringá, PR., Brazil. Phone: +5544 3011-4809, Fax: +5544 3011-4860, E-mail: tiesvidzinski@uem.br or terezinha.svidzinski@gmail.com

<http://dx.doi.org/10.17179/excli2019-1672>

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>).



Supplementary Figure 1: Phylogenetic analysis of *Rhodotorula mucilaginosa*, based on ITS sequences constructed with Maximum likelihood, based on the Tamura-Nei model + Gamma distribution (T92+G) implemented in MEGA v.7. Bootstrap support was calculated from 1000 replicates. (T) = type strain of the species. Bootstrap values > 80 % were considered statistically significant.