









Supplementary material to:

Original article:

**PLASMA AND URINARY CONCENTRATIONS OF
ARACHIDONIC ACID-DERIVED EICOSANOIDS ARE ASSOCIATED
WITH DIABETIC KIDNEY DISEASE**

Sonia Mota-Zamorano¹, Nicolás R. Robles², Juan Lopez-Gomez³, Bárbara Cancho²,
Luz M. González¹, Guadalupe Garcia-Pino⁴, María Luisa Navarro-Pérez⁵,
Guillermo Gervasini^{1*}

¹ Department of Medical and Surgical Therapeutics, Division of Pharmacology,
Medical School, University of Extremadura, Badajoz, Spain

² Service of Nephrology, Badajoz University Hospital, Badajoz, Spain

³ Service of Clinical Analyses, Badajoz University Hospital, Badajoz, Spain


⁴ Service of Nephrology, Zafra Hospital, Badajoz, Spain

⁵ Department of Biomedical Sciences, University of Extremadura, Badajoz, Spain


* **Corresponding author:** Guillermo Gervasini, PhD, Department of Medical and Surgical
Therapeutics, Division of Pharmacology, Medical School, University of Extremadura,
Av. Elvas s/n 06071, Badajoz, Spain. Phone: +34 924 289 824, E-mail: ggervasi@unex.es

<http://dx.doi.org/10.17179/excli2021-3408>


This is an Open Access article distributed under the terms of the Creative Commons Attribution License
(<http://creativecommons.org/licenses/by/4.0/>).



Supplementary Figure 1: Chromatogram showing retention times for the araquidonic-acid metabolites in plasma



Supplementary Figure 2: Distribution of 14,15- and 11,12-DHET plasma levels according to albuminuria in diabetic and non-diabetic subjects



Supplementary Figure 3: Distribution of 20-HETE plasma levels according to abuminuria (A), glomerular filtration rate (eGFR) (B), and association with diabetic nephropathy (C)