

Case series report:

MANUAL THERAPY AND EXERCISE FOR PATIENTS WITH CLUSTER HEADACHE

Lucía de-la-Puente-Ranea^{1,6}, Alfonso Gil-Martínez^{1,2,4,6*} , Oscar Rodríguez-Lopez^{5,6} , Pablo González-Gutiérrez¹, María Ángeles Mangas-Guijarro³, Gonzalo Navarro-Fernández^{1,2,6} 

- ¹ Departamento de Fisioterapia, Centro Superior de Estudios Universitarios La Salle, Universidad Autónoma de Madrid, Madrid (Spain)
- ² CranioSPain Research Group, Centro Superior de Estudios Universitarios La Salle, Madrid (Spain)
- ³ Servicio de Neurología, Hospital Universitario La Paz, Madrid, Spain
- ⁴ Unidad de Fisioterapia, Hospital Universitario La Paz (IdiPAZ), Madrid, Spain
- ⁵ Departamento de Fisioterapia, CEU San Pablo, Madrid, Spain
- ⁶ Instituto de Rehabilitación Funcional, Centro Superior de Estudios Universitarios La Salle, Madrid (Spain)

* **Corresponding author:** Alfonso Gil-Martínez, Unidad de Fisioterapia, Hospital Universitario La Paz. Paseo de la Castellana, 261. 28046. Madrid. Spain; Tel.: + 0034 666137908; E-mail: fongilmar@gmail.com

<http://dx.doi.org/10.17179/excli2021-3763>

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>).

ABSTRACT

The aim of this case series is to clarify if a physiotherapy program can reduce the frequency, intensity and duration of the headache episodes in patients with cluster headache. A 7-case series with cluster headache patients was conducted. Every subject received physiotherapy treatment based on manual therapy and exercise, maintaining pharmacological treatment prescribed by the neurologist. Frequency, intensity and duration of the episodes, pressure pain thresholds (PPT) and psychological variables were measured 5 times: pre-intervention, post-intervention, 1 month follow-up, 2 months follow-up and 3 months follow-up. Two of the seven subjects decreased their frequency of headaches over 50 % and another in 16,67 %. There were no significant changes in duration or intensity. Results also showed an improvement in impact of headache in 6 of 7 cases. Those cases that decreased more their headache frequency also decreased their pain catastrophizing. A program of physiotherapy based in manual therapy and exercise, might be an effective and safe complement to decrease the frequency of the episodes of CH in short-term (4 months follow-up) including interdisciplinary working with neurologists and other health care professionals.

Keywords: Disability, manual therapy, exercise, cluster headache, pain, physiotherapy

Abbreviations:

CH	Cluster Headache	PCS	Pain Catastrophizing Scale
CCH	Chronic Cluster Headache	PA	Postero-Anterior
ECH	Episodic Cluster Headache	C2	Second Cervical Vertebra
PPT	Pressure Pain Threshold	V1	First Trigeminal Branch
HIT6	Headache Impact Test	V2	Second Trigeminal Branch

INTRODUCTION

Cluster headache (CH) is the most disabling type of primary headache and it has been defined as a trigeminal-autonomous headache subtype by the International Classification of Headache Disorders, characterized by the onset of high intensity and short duration episodes of unilateral pain associated with autonomic symptoms (Halker et al., 2010). Pain usually comprises the orbital, supraorbital and temporal areas, starting frequently in the upper jaw (May, 2013). In terms of frequency, CH presents both chronic (CCH) and episodic forms (ECH) (Chaibi and Russell, 2014) and it is more usual in men (3:1), with an age of onset between twenty and forty years old (Weaver-Agostoni, 2013).

Based on clinical features of CH, it has been proposed in some studies that CH patients could have signs and symptoms suggesting a central sensitization process. It has been showed that CH patients have hyperalgesia (Bono et al., 1996; Ladda et al., 2006; Fernández-de-Las-Peñas et al., 2011), allodynia (Marmura et al., 2009), high risk of depression (Liang et al., 2013) and decreased health-related quality of life even when appropriate treatments are used (D'Amico et al., 2002). In 2019, Gil-Martínez et al. published a study determining that CH patients showed cranial and extracranial hyperalgesia, high levels of impact of headache, higher levels of anxiety and depression and lower levels of quality of life when compared with healthy subjects. Even more interesting is the fact that anxiety and depression were positively correlated with duration and frequency of headache, respectively (Gil-Martínez et al., 2019).

Regarding this complex clinical situation and the burden of the disease (Jensen et al. 2007), some pharmacological and non-pharmacological treatments have been proposed in order to reduce the impact of the headache in patient's lives. Among pharmacological options, verapamil, topiramate and lithium stand out as a preventive treatment and oxygen 100 %, subcutaneous sumatriptan, oral corticosteroids and occipital nerve block as abortive treatment (Becker, 2013).

Furthermore, other conservative treatment frequently used in the management of primary headache patients is physical therapy and therapeutic exercise. However, although the efficacy of physiotherapy has been studied for other primary headaches (Chaibi et al., 2011), there are no studies evaluating the effects of physiotherapy in patients with CH controlled with basal medication. There is only one study published in relation to physical therapy on CH, a case report in which manual therapy and therapeutic exercise, combined with neurostimulation and medication, seems to be useful in the treatment of CH. Regarding all primary headaches, some of the techniques which have demonstrated hypoalgesic effects are manual therapy (Chaibi and Russell, 2014) and therapeutic exercise (Gil-Martínez et al., 2013). Specifically, it appears that the upper neck region may be an important target for physiotherapeutic approaches in CH patients. Recently, an article was published showing promising results in occipital nerve stimulation (Díaz-de-Terán et al., 2021).

Therefore, the aim of this case series is to clarify if a physiotherapy program based on manual therapy and therapeutic exercise can reduce the frequency, intensity and duration of the headache episodes in patients with CH.

METHODS

This study was carried out in a private physiotherapy center and in a tertiary public hospital. The study had a total duration of 8 months.

Participants and evaluation description

Cases were adults aged 18 to 65 years who had been diagnosed with CCH or ECH by a neurologist. All patients signed the informed consent before the beginning of the study. All subjects with systemic diseases, fibromyalgia, peripheral neuropathies, craneomandibular dysfunctions and recent trauma or surgical interventions in neck or facial areas were excluded.

Five evaluations were included in the case series: pre-intervention, post-intervention and

one, two and three months after the post-intervention. In the first evaluation, the subjects who accepted the participation in the study were informed about the study by a physiotherapist. After that, a personal interview was carried out to obtain information about the pharmacological treatment, demographical data and specific characteristics of pain, such as frequency, intensity and duration of the headache.

The pain pressure threshold (PPT) of facial and peripheral areas was measured in every evaluation using a digital algometer (Fx.25 Force Gage model, Wagner Instruments, Greenwich, CT, USA). The PPT bilateral measurements were: the two upper branches of the trigeminal nerve (V1, in supraorbital keyhole and V2, in infraorbital keyhole) and tibialis anterior. To conclude, each subject filled the Headache Impact Test-6 (HIT6) (Sauro et al., 2010), the Pain Catastrophizing Scale (PCS) (Olmedilla Zafra et al., 2013) and the disability and daily life affection were measured with a non-verbal scale where 0 was no affection and 10 the greatest affection.

Treatment

A physiotherapy program was applied, without making any changes in basal medication prescribed by the neurologist. Eight sessions were carried out; organized in 2 sessions per week for 1 month. The first technique was a postero-anterior (PA) vertebral mobilization on the second cervical vertebra (C2) at 0.5 Hz during 3 series of 2 minutes, resting 30 seconds between each series (La Touche et al., 2013). The second technique was a neural mobilization of the trigeminal nerve, organized in 30 repetitions at 0.5 Hz, making 10 repetitions globally and 10 repetitions each side, increasing the tension in the auriculotemporal nerve area. After that deep flexor motor control home exercises were prescribed as described by Harris et al. (2005) (Figure 1).

RESULTS

Seven subjects were recruited (2 with CCH and 5 with ECH). Among the patients with ECH, 4 of them were in remission period (number 4 to number 7) and only 1 subject was in active period (case number 3). Demographic characteristics are described in Table 1.




Figure 1: Physiotherapy program

Table 1: Demographic characteristics of the patients

N	7
Age [years (SD)]	43.29 (5.12)
Gender [men/women]	5/2
Height [cm (SD)]	174.43 (4.28)
Weight [Kg (SD)]	71.09 (2.99)
BMI [mean (SD)]	23.44 (1.00)

SD, Standard deviation; BMI, Body mass index

One of the two subjects with CCH decreased its headache frequency more than a 50 % of his basal frequency after treatment; and the other subject decreased his headache frequency a 16.67 %. Among the patients with ECH, only one of the cases in active period decreased the headache frequency in more than a 50 %. There were no changes in ECH in remission period (Table 2).

Concerning PPT, an increase in V1 mean from pre-treatment to 3rd month follow-up

was observed, from 0.53 (0.21) to 0.63 (0.27). A decrease was also reported in HIT-6 mean from pre-treatment to 3rd month follow-up, from 60.43 (8.85) to 50.57 (8.16), and in PCS from 32.86 (3.17) to 21.14 (18.18) (Tables 3 and 4).

Disability and daily life affection were also decreased in patients with CCH, and a complete disability reduction was reported in the subject with ECH in active period, maintaining with no differences in subjects with ECH in remission period.

DISCUSSION

According to clinical guidelines in CH, there are differences between patients with CCH and ECH (Lademann et al., 2015). Nevertheless, in this case series patients from both categories have been included to approach the epidemiology of both entities.

Considering the obtained data, a decrease in frequency of headaches was reported in subjects with both CCH and ECH in active period. These results may have been caused by the analgesic effect generated by the phy-

Table 2: Subjective evolution of pain per case

CASES	Variables	PRE	POST	1 MONTH	2 MONTHS	3 MONTHS
1	Freq.	60	35	4	-	20
	Dur.	60	17.50	7.50	-	30
	Int.	10	9.50	9	-	8
2	Freq.	30	30	14	30	25
	Dur.	90	90	52	60	90
	Int.	8	6	6	7	6
3	Freq.	30	10	0	-	0
	Dur.	20	15	0	-	0
	Int.	9	5	0	-	0
4	Freq.	0	2*	0	1*	0
	Dur.	0	12.50*	0	10*	0
	Int.	0	2*	0	4*	0
5	Freq.	0	0	0	0	0
	Dur.	0	0	0	0	0
	Int.	0	0	0	0	0
6	Freq.	0	0	0	0	0
	Dur.	0	0	0	0	0
	Int.	0	0	0	0	0
7	Freq.	0	0	-	0	0
	Dur.	0	0	-	0	0
	Int.	1.5	0	-	1	0

Freq, Frequency (days); Dur, Duration (minutes); Int, Intensity (0 to 10).

*No headache was reported, but aura was.

Table 3: Somatosensorial characteristics of the affected side and psychological characteristics

	CASES	PPT V1	PPT V2	PPT TIBIALIS ANT	HIT-6	PCS
1	Pre-Post	0.61 - 0.74	1.58 - 1.00	3.12 - 3.22	76 - 70	50 - 47
	Post-3month	0.74 - 0.55	1.00 - 1.25	3.22 - 2.91	70 - 56* †	47 - 05*
	Pre-3month	0.61 - 0.55	1.58 - 1.25	3.12 - 2.91	76 - 56* †	50 - 05*
2	Pre-Post	0.27 - 0.45	0.41 - 0.58	1.97 - 1.24	67 - 58* †	18 - 05*
	Post-3month	0.45 - 0.46	0.58 - 0.46	1.24 - 1.28	58 - 60	05 - 17*
	Pre-3month	0.27 - 0.46	0.41 - 0.46	1.97 - 1.28	67 - 60	18 - 17
3	Pre-Post	0.47 - 0.64	0.60 - 1.10	1.80 - 3.48*	60 - 42* †	26 - 14*
	Post-3month	0.64 - 1.14	1.10 - 1.75	3.48 - 5.22*	42 - 36	14 - 00*
	Pre-3month	0.47 - 1.14	0.60 - 1.75	1.80 - 5.22*	60 - 36* †	26 - 00*
4	Pre-Post	0.47 - 0.68	0.47 - 0.70	1.34 - 1.29	50 - 63* †	34 - 27
	Post-3month	0.68 - 0.62	0.70 - 0.55	1.29 - 1.02	63 - 46* †	27 - 23
	Pre-3month	0.47 - 0.62	0.47 - 0.55	1.34 - 1.02	50 - 46	34 - 23*
5	Pre-Post	0.93 - 0.45	1.69 - 0.80	6.07 - 3.53*	59 - 59	17 - 11
	Post-3month	0.45 - 0.49	0.80 - 0.59	3.53 - 2.19	59 - 51*	11 - 14
	Pre-3month	0.93 - 0.49	1.69 - 0.59	6.07 - 2.19*	59 - 51*	17 - 14
6	Pre-Post	0.53 - 0.78	0.60 - 0.94	2.21 - 2.86	59 - 50* †	38 - 42
	Post-3month	0.78 - 0.81	0.9 - 0.47	2.86 - 1.52	50 - 48	42 - 37
	Pre-3month	0.53 - 0.81	0.60 - 0.47	2.21 - 1.52	59 - 48* †	38 - 37
7	Pre-Post	0.40 - 0.38	0.62 - 0.39	2.86 - 1.53	52 - 53	47 - 50
	Post-3month	0.38 - 0.33	0.39 - 0.47	1.53 - 1.46	53 - 57	50 - 52
	Pre-3month	0.40 - 0.33	0.62 - 0.47	2.86 - 1.46	52 - 57	47 - 52

PPT, Pressure Pain Threshold; V1, first trigeminal branch of the trigeminus nerve; V2, second trigeminal branch of the trigeminus nerve; Tibialis Ant, Tibialis anterior

† Clinically Relevant Change HIT-6 = >8

* Minimal Detectable Change HIT-6 = 8

* Minimal Detectable Change PCS = 9

* Minimal Detectable Change Tibialis Anterior 1.57

siotherapy techniques, which in addition to the pharmacological treatment may have this effect. Also, the results obtained in this case series were similar to those obtained in another previous study (Navarro-Fernández et al., 2019).

Joint mobilization has demonstrated an analgesic effect in other types of headache (Chaibi et al., 2011; La Touche et al., 2013), activating descending pain pathways such as noradrenergic and serotonergic pathways,

which involve both supraspinal and grey periaqueductal substance areas (Skyba et al., 2003).

Furthermore, it has been reported that neural mobilization produced a normalization in the expression of astrocytes and microglial cells in the posterior horn (Martins et al., 2011; Santos et al., 2012). Thus, neural mobilization could decrease glial activity, which is implied in chronification of pain.

Table 4: Means and Standard deviation of somatosensorial characteristics of the affected side and psychological characteristics (all cases)

		V1	V2	TIBIALIS ANT	HIT-6	PCS
Mean (SD)	Pre	0.53 (0.21)	0.85 (0.54)	2.77 (1.58)	60.43 (8.85)	32.86 (3.17)
	Post	0.60 (0.16)	0.79 (0.25)	2.45 (1.05)	56.43 (9.11)	28.00 (18.51)
	3 months	0.63 (0.27)	0.79 (0.50)	2.22 (1.46)	50.57 (8.16)	21.14 (18.18)

PPT, Pressure Pain Threshold; V, first trigeminal branch of the trigeminal nerve; V2, second trigeminal branch of the trigeminal nerve; Tibialis Ant, Tibialis anterior; SD, Standard Deviation

Likewise, it has been observed that a neck and cranium motor control training can reduce frequency, intensity and duration of headaches in the long term (Busch and Gaul, 2008). This could be due to an activation in opioid pathways after exercising, which triggers a liberation of β -endorphins in hypothalamus and grey periaqueductal substance areas, activating pain inhibition pathways. Moreover, there are reviews which support a treatment based in therapeutic exercise with cardiovascular exercise in primary headaches (Gil-Martínez et al., 2013), even in a preventive treatment (Varkey et al., 2011).

No moderate changes in the intensity, duration, and number of episodes of CH were observed in subjects in the active period; hence, the need for future studies with a larger sample size.

Regarding PPT, an increase in the mean of V1 when comparing pre-intervention and 3 months follow-up and a decrease in V2 and tibialis anterior was reported. These last results are contrary to previous studies, in which the combination of the applied techniques in the cervical segment triggered an increase of the PPT in facial areas (La Touche et al., 2009).

Nevertheless, the reported changes in tibialis anterior do not reach the minimal detectable change (Walton et al., 2011), which indicates that the sample size should be increased so that the effects of the physiotherapy techniques on PPT in CH are observed.

With reference to psychological factors, 6 out of 7 cases improved the daily life affection of the CH. This may be a future line of research, as high results in this variable have been reported previously both in CCH and

ECH in active or remission period (Torkamani et al., 2015).

The obtained results indicate that those cases with a higher rate of episodes decrease more their catastrophism. This may indicate a relationship between pain catastrophizing and other headache-related variables such as frequency, chronicity, duration, self-sufficiency (Bond et al., 2015), disability and pain intensity (Queralto, 2005). Such relationship has been already proved in the classic model of Vlaeyen and Linton (2000).

In a similar way, subjects who decreased their rate of episodes also decreased their disability. Similar results were reported by Ragi et al. in a study in which the direct contact between patient and professional triggered a reduction in disability in patients with migraine (Raggi et al., 2012). No adverse effects were reported.

Implications for physiotherapy practice

The results obtained in this case series seem to be supportive of the idea that physical therapy interventions based on exercise and manual therapy could be beneficial for cluster headache patients. However, considering that this is a case series, these preliminary results should be interpreted with caution.

Exercise and manual therapy reduced disability and daily live affection of cluster headache patients in this case series.

Physical therapy interventions based on manual therapy and exercise seemed to improve physical and psychological variables of cluster headache patients.

Limitations

There are many limitations to consider in this case series. First, the results cannot be extrapolated to the rest of population. Nevertheless, the investigators consider that this case of series is a preliminary tool to further investigations. Second, the data interpretation should be conservative, due to the study design and the lack of a control group. Third, the consumption of analgesic drugs may have influenced in the patient condition. It should also be considered that the recruitment was partially carried out in an Association of CH, which might imply common expectation on the treatment. The lack of literature related with CH indicates a need to carry out more investigations that observe the effect of physiotherapy on CH, making a separated classification of CCH, ECH in both active and remission period. Last, the lack of previous studies in which PPTs minimum detectable change and clinically relevant change were obtained in the facial area.

In conclusion, a program of physiotherapy based in manual therapy and exercise, might be an effective and safe complement to decrease the frequency of the episodes of CH in short-term (4 months follow-up) including interdisciplinary working with neurologists and other health care professionals.

Conflict of interest

No potential conflict of interest relevant to this article was reported. This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Acknowledgments

The authors would like to sincerely thank the patients who graciously participated in this case of series. We also thank Centro de Estudios Superiores La Salle for its assistance with the English edition of the manuscript.

REFERENCES

Becker WJ. Cluster headache: Conventional pharmacological management. *Headache*. 2013;53: 1191–6.

Bond DS, Buse DC, Lipton RB, Thomas JG, Rathier L, Roth J, et al. Clinical pain catastrophizing in women with migraine and obesity. *Headache*. 2015;55:923–33.

Bono G, Antonaci F, Sandrini G, Pucci E, Rossi F, Nappi G. Pain pressure threshold in cluster headache patients. *Cephalalgia*. 1996;16:62–6; Discussion 3-4.

Busch V, Gaul C. Exercise in migraine therapy--is there any evidence for efficacy? A critical review. *Headache*. 2008; 48:890–9.

Chaibi A, Russell MB. Manual therapies for primary chronic headaches: a systematic review of randomized controlled trials. *J Headache Pain*. 2014;15:67.

Chaibi A, Tuchin PJ, Russell MB. Manual therapies for migraine: a systematic review. *J Headache Pain*. 2011; 12:127–33.

D'Amico D, Rigamonti A, Solari A, Leone M, Usai S, Grazi L, et al. Health-related quality of life in patients with cluster headache during active periods. *Cephalalgia*. 2002;22:818–21.

Díaz-de-Terán J, Membrilla JA, Paz-Solís J, de Lorenzo I, Roa J, Lara-Lara M, et al. Occipital nerve stimulation for pain modulation in drug-resistant chronic cluster headache. *Brain Sci*. 2021;11:236.

Fernández-de-Las-Peñas C, Ortega-Santiago R, Cuadrado ML, López-de-Silanes C, Pareja JA. Bilateral widespread mechanical pain hypersensitivity as sign of central sensitization in patients with cluster headache. *Headache*. 2011;51:384–91.

Gil-Martínez A, Kindelan-Calvo P, Agudo-Carmona D, Muñoz-Plata R, López-de-Uralde-Villanueva I, et al. Therapeutic exercise as treatment for migraine and tension-type headaches: A systematic review of randomised clinical trials. *Rev Neurol*. 2013;57:433–43.

Gil-Martínez A, Navarro-Fernández G, Mangas-Guijarro MÁ, Díaz-de-Terán J. Hyperalgesia and central sensitization signs in patients with cluster headache: A cross-sectional study. *Pain Med*. 2019;20: 2562–70.

Halker R, Vargas B, Dodick DW. Cluster headache: diagnosis and treatment. *Semin Neurol*. 2010;30:175–85.

Harris KD, Heer DM, Roy TC, Santos DM, Whitman JM, Wainner RS. Reliability of a measurement of neck flexor muscle endurance. *Phys Ther*. 2005;85:1349–55.

Jensen RM, Lyngberg A, Jensen RH. Burden of cluster headache. *Cephalalgia*. 2007;27:535–41.

- La Touche R, Fernández-de-las-Peñas C, Fernández-Carnero J, Escalante K, Angulo-Díaz-Parreño S, Paris-Alemany A, et al. The effects of manual therapy and exercise directed at the cervical spine on pain and pressure pain sensitivity in patients with myofascial temporomandibular disorders. *J Oral Rehabil.* 2009;36:644–52.
- La Touche R, Paris-Alemany A, Mannheimer JS, Angulo-Díaz-Parreño S, Bishop MD, López-Valverde-Centeno A, et al. Does mobilization of the upper cervical spine affect pain sensitivity and autonomic nervous system function in patients with cervicocraniofacial pain? A randomized-controlled trial. *Clin J Pain.* 2013;29:205–15.
- Ladda J, Straube A, Förderreuther S, Krause P, Eggert T. Quantitative sensory testing in cluster headache: increased sensory thresholds. *Cephalalgia.* 2006;26:1043–50.
- Lademann V, Jansen J-P, Evers S, Frese A. Evaluation of guideline-adherent treatment in cluster headache. *Cephalalgia.* 2015;36:760–4.
- Liang J-F, Chen Y-T, Fuh J-L, Li S-Y, Liu C-J, Chen T-J, et al. Cluster headache is associated with an increased risk of depression: a nationwide population-based cohort study. *Cephalalgia.* 2013;33:182–9.
- Marmura MJ, Abbas M, Ashkenazi A. Dynamic mechanical (brush) allodynia in cluster headache: a prevalence study in a tertiary headache clinic. *J Headache Pain.* 2009;10:255–8.
- Martins DF, Mazzardo-Martins L, Gadotti VM, Nascimento FP, Lima DAN, Speckhann B, et al. Ankle joint mobilization reduces axonotmesis-induced neuropathic pain and glial activation in the spinal cord and enhances nerve regeneration in rats. *Pain.* 2011;152:2653–61.
- May A. Diagnosis and clinical features of trigemino-autonomic headaches. *Headache.* 2013;53:1470–8.
- Navarro-Fernández G, de-la-Puente-Ranea L, Gandía-González M, Gil-Martínez A. Endogenous neurostimulation and physiotherapy in cluster headache: A clinical case. *Brain Sci.* 2019;9:60.
- Olmedilla Zafra A, Ortega Toro E, Cano LA. Validation of the pain catastrophizing scale in Spanish athletes. *Cuad Psicol del Deport.* 2013;13:83–93.
- Queralto JM. Análisis de los factores psicológicos moduladores del dolor crónico benigno. *Anu Psicol.* 2005;36:37–60.
- Raggi A, Giovannetti AM, Quintas R, D’Amico D, Cieza A, Sabariego C, et al. A systematic review of the psychosocial difficulties relevant to patients with migraine. *J Headache Pain.* 2012;13:595–606.
- Santos FM, Silva JT, Giardini AC, Rocha PA, Achermann APP, Alves AS, et al. Neural mobilization reverses behavioral and cellular changes that characterize neuropathic pain in rats. *Mol Pain.* 2012;8:57.
- Sauro KM, Rose MS, Becker WJ, Christie SN, Giammarco R, Mackie GF, et al. HIT-6 and MIDAS as measures of headache disability in a headache referral population. *Headache.* 2010;50:383–95.
- Skyba DA, Radhakrishnan R, Rohlwing JJ, Wright A, Sluka KA. Joint manipulation reduces hyperalgesia by activation of monoamine receptors but not opioid or GABA receptors in the spinal cord. *Pain.* 2003;106:159–68.
- Torkamani M, Ernst L, Cheung LS, Lambru G, Matharu M, Jahanshahi M. The neuropsychology of cluster headache: cognition, mood, disability, and quality of life of patients with chronic and episodic cluster headache. *Headache.* 2015;55:287–300.
- Varkey E, Cider A, Carlsson J, Linde M. Exercise as migraine prophylaxis: A randomized study using relaxation and topiramate as controls. *Cephalalgia.* 2011;31:1428–38.
- Vlaeyen JW, Linton SJ. Fear-avoidance and its consequences in chronic musculoskeletal pain: A state of the art. *Pain.* 2000;85:317–32.
- Walton DM, Macdermid JC, Nielson W, Teasell RW, Chiasson M, Brown L. Reliability, standard error, and minimum detectable change of clinical pressure pain threshold testing in people with and without acute neck pain. *J Orthop Sports Phys Ther.* 2011;41:644–50.
- Weaver-Agostoni J. Cluster headache. *Am Fam Physician.* 2013;88:122–8.