

EDITORIAL:

ANALYSIS OF REACTIVE OXYGEN SPECIES

Hermann M. Bolt¹, Jan G. Hengstler¹, Joanna Stewart¹ ✉

1 Leibniz-Institut für Arbeitsforschung an der TU Dortmund, Leibniz Research Centre for Working Environment and Human Factors (*IfADO*), Ardeystrasse 67, 44139 Dortmund, Germany

✉ Corresponding author: e-mail: stewart@ifado.de

Reactive oxygen species play key roles as signalling molecules (Thannickal et al., 2000; Landar and Darley-Usmar, 2003; Pryor et al., 2006; Hengstler and Bolt, 2008) and are also involved in many pathogenic processes (Cederbaum et al., 2009; Sebai et al., 2009). Oxidative stress induction is a well-documented mechanism of many non-genotoxic carcinogens (Mátes et al., 2008; Hengstler and Bolt, 2007; Schug et al., 2008; Dewa et al., 2009; Binner et al., 2008). Particularly metal toxicity and carcinogenicity often involves generation of reactive oxygen species as a key mechanism (Beyersmann and Hartwig, 2008; Wang et al., 2009; Cervinková et al., 2009; Bolt and Hengstler, 2008; Verstraeten et al., 2008; Glahn et al., 2008; Pardo Andreu et al., 2009; Arivarasu et al., 2008).

Given the importance of oxidative stress numerous methods have been established and optimized to reliably quantify the levels of reactive oxygen species in tissues and cells (Dikalov et al., 2007; Borgmann 2009; Setsukinai et al., 2003; Yao et al., 2004; Tarpey and Fridovich, 2001; Tarpey et al., 2004). Some of the most frequently applied techniques include:

1. *Detection of 2-hydroxyethidium formation from dihydroethidium*

Dihydroethidium reacts with O_2^- to form 2-hydroxyethidium which can be quantified by HPLC (Zhao et al., 2003, 2005; Fink et al., 2004; Dikalov et al., 2007).

2. *DCFH-DA as a fluorescent probe to detect intracellular H_2O_2*

DCFH-DA (2',7'-dichlorofluorescein diacetate) can pass cell membranes and is cleaved by intracellular esterases to 2',7'-dichlorofluorescein (DCFH) and thereby trapped within the cells. A variety of reactive oxygen species oxidize DCFH to the fluorescent DCF (2',7'-dichlorofluorescein) (Bass et al., 1983; Khatri et al., 2004; Liu et al., 2003; Seshiah et al., 2002; Abdalla et al., 2005; Marchesi et al., 1999; Rota et al., 1999a, b; Wrona and Wardman, 2006; Kim et al., 2006). However, DCFH is easily autoxidized, resulting in unspecific fluorescence upon exposure to light (Hempel et al., 1999; Setsukinai et al., 2003). Therefore, research is ongoing to design more specific fluorescence probes (Setsukinai et al., 2003).

3. *Amplex red for detection of extracellular H_2O_2*

Amplex red (N-acetyl-3,7-dihydroxyphen-oxazine) is a nonfluorescent molecule that is oxidized to the fluorescent resorufin by horseradish peroxidase in the presence of H_2O_2 (Zhou et al., 1997). The amplex red technique can be applied to quantify H_2O_2 released by cells and tissues in cell-free supernatants.

4. *Cytochrome c reduction to detect O₂⁻*

Ferricytochrome c is reduced to ferrocyanochrome c in the presence of O₂⁻ (Landmesser et al., 2003; Guzik et al., 2002; Gongora et al., 2006). Ferrocyanochrome c shows a different absorption spectrum compared to ferricytochrome and can easily be quantified spectrophotometrically. However, sensitivity of this technique is limited.

5. *Chemiluminescence based methods*

Exposure to chemiluminescent probes, such as lucigenin, to O₂⁻ leads to release of a photon from the probe, which can be detected luminometrically or by a scintillation counter (Li et al., 1998; Faulkner and Fridovich, 1993; Daiber et al., 2004). However, recently also rapid and easy to handle thermoluminescence based techniques have been established, where no extraction procedures are needed (Schumann et al., 2009). Using directly collected cell material the CCl₄ induced increase in thermoluminescence high temperature bands correlated well with the results of established biochemical methods.

6. *Electron spin resonance (ESR) and electrochemical techniques*

Free radicals may cause absorption of microwave radiation (Dikalov et al., 2007). Based on this principle ESR is a sensitive technique for studies of tissues. However, the technique is expensive and requires trained experts. Similarly, electrochemical methods where electrodes are applied to measure for instance H₂O₂ are technically challenging (Borgmann, 2009). However, an advantage of electrochemical methods is the possibility of real time quantification of reactive oxygen species in the microenvironment of cells. Recently, Tangkosakul et al. have presented a new technique for antioxidant activity in this years edition of the EXCLI J (Tangkosakul et al., 2009). This technique is based on lactate dehydrogenase oxidation and may find a place in the specific field of screening for compounds with antioxidative capacity. Presently, numerous studies are performed aimed at identifica-

tion of compounds with antioxidative properties (Yang et al., 2009; Yu et al., 2009; Adam and Laufs, 2008; Helal and Helal, 2009). For this purpose easy to handle, quantitative, reproducible and cheap techniques are required that also allow high throughput analysis. All these requirements are fulfilled by the new LDH-based technique of Tangkosakul et al. (2009) which therefore may become a frequently applied method in screening for compounds with antioxidative properties.

REFERENCES

- Abdalla MY, Ahmad IM, Spitz DR, Schmidt WN, Britigan BE. Hepatitis C virus-core and non structural proteins lead to different effects on cellular antioxidant defenses. *J Med Virol* 2005;76:489-97.
- Adam O, Laufs U. Antioxidative effects of statins. *Arch Toxicol* 2008;82:885-92.
- Arivarasu NA, Fatima S, Mahmood R. Oral administration of potassium dichromate inhibits brush border membrane enzymes and alters anti-oxidant status of rat intestine. *Arch Toxicol* 2008;82:951-8.
- Bass DA, Parce JW, Dechatelet LR, Szejda P, Seeds MC, Thomas M. Flow cytometric studies of oxidative product formation by neutrophils: a graded response to membrane stimulation. *J Immunol* 1983;130:1910-7.
- Beyersmann D, Hartwig A. Carcinogenic metal compounds: recent insight into molecular and cellular mechanisms. *Arch Toxicol* 2008;82:493-512.
- Binner C, Selinski S, Barysch MJ, Pölcher C, Schormann W, Hermes M, Brulport M, Bauer A, Rudolph C, Bedawy E, Schug M, Golka K, Hasenclever D, Trauer H, Lessig R, Bolt HM, Ickstadt K, Hengstler JG. Munich Oktoberfest experience: remarkable impact of sex and age in ethanol intoxication. *Arch Toxicol* 2008;82:933-9.

- Bolt HM, Hengstler JG. Aluminium and lead toxicity revisited: mechanisms explaining the particular sensitivity of the brain to oxidative damage. *Arch Toxicol* 2008;82:787-8.
- Borgmann S. Electrochemical quantification of reactive oxygen and nitrogen: challenges and opportunities. *Anal Bioanal Chem* 2009;394:95-105.
- Cederbaum AI, Lu Y, Wu D. Role of oxidative stress in alcohol-induced liver injury. *Arch Toxicol* 2009;83:519-48.
- Cervinková Z, Kriváková P, Lábajová A, Rousar T, Lotková H, Kucera O, Endlicher R, Cervinka M, Drahotka Z. Mechanisms participating in oxidative damage of isolated rat hepatocytes. *Arch Toxicol* 2009;83:363-72.
- Daiber A, Oelze M, August M, Wendt M, Sydow K, Wieboldt H, Kleschyov AL, Munzel T. Detection of superoxide and peroxynitrite in model systems and mitochondria by the luminol analogue L-012. *Free Radic Res* 2004;38:259-69.
- Dewa Y, Nishimura J, Muguruma M, Jin M, Kawai M, Saegusa Y, Okamura T, Umemura T, Mitsumori K. Involvement of oxidative stress in hepatocellular tumor-promoting activity of oxfendazole in rats. *Arch Toxicol* 2009;83:503-11.
- Dikalov S, Griendling KK, Harrison DG. Measurement of reactive oxygen species in cardiovascular studies. *Hypertension* 2007;49:717-27.
- Faulkner K, Fridovich I. Luminol and lucigenin as detectors for O₂·-. *Free Radic Biol Med* 1993;15:447-51.
- Fink B, Laude K, McCann L, Doughan A, Harrison DG, Dikalov S. Detection of intracellular superoxide formation in endothelial cells and intact tissues using dihydroethidium and an HPLC-based assay. *Am J Physiol* 2004;287:C895-902.
- Glahn F, Schmidt-Heck W, Zellmer S, Guthke R, Wiese J, Golka K, Hergenröder R, Degen GH, Lehmann T, Hermes M, Schormann W, Brulport M, Bauer A, Bedawy E, Gebhardt R, Hengstler JG, Foth H. Cadmium, cobalt and lead cause stress response, cell cycle deregulation and increased steroid as well as xenobiotic metabolism in primary normal human bronchial epithelial cells which is coordinated by at least nine transcription factors. *Arch Toxicol* 2008;82:513-24.
- Gongora MC, Qin Z, Laude K, Kim HW, McCann L, Folz JR, Dikalov S, Fukai T, Harrison DG. Role of extracellular superoxide dismutase in hypertension. *Hypertension* 2006;48:473-81.
- Guzik TJ, Mussa S, Gastaldi D, Sadowski J, Ratnatunga C, Pillai R, Channon KM. Mechanisms of increased vascular superoxide production in human diabetes mellitus: role of NAD(P)H oxidase and endothelial nitric oxide synthase. *Circulation* 2002;105:1656-62.
- Helal GK, Helal OK. Metallothionein attenuates carmustine-induced oxidative stress and protects against pulmonary fibrosis in rats. *Arch Toxicol* 2009;83:87-94.
- Hempel SL, Buettner GR, O'Malley YQ, Wessels DA, Flaherty DM. Dihydrofluorescein diacetate is superior for detecting intracellular oxidants: comparison with 2',7'-dichlorodihydrofluorescein diacetate, 5(and 6)-carboxy-2',7'-dichlorodihydrofluorescein diacetate, and dihydrorhodamine 123. *Free Radic Biol Med* 1999;27:146-59.
- Hengstler JG, Bolt HM. Induction and control of oxidative stress. (Editorial). *Arch Toxicol* 2007;81:823-4.
- Hengstler JG, Bolt HM. Oxidative stress: from modification of cell-cycle related events, secondary messenger function, dysregulation of small GTPases, protein kinases and phosphatases to redox-sensitive cancer models. *Arch Toxicol* 2008;82:273-99.

Khatri JJ, Johnson C, Magid R, Lessner SM, Laude KM, Dikalov SI, Harrison DG, Sung HJ, Rong Y, Galis ZS. Vascular oxidant stress enhances progression and angiogenesis of experimental atheroma. *Circulation* 2004;109:520-5.

Kim YM, Lim JM, Kim BC, Han S. Cu,Zn-superoxide dismutase is an intracellular catalyst for the H₂O₂-dependent oxidation of dichlorodihydrofluorescein. *Mol Cells* 2006;21:161-5.

Landar A, Darley-Usmar VM. Nitric oxide and cell signaling; modification of redox tone and protein modification. *Amino Acids* 2003;25:313-21.

Landmesser U, Dikalov S, Price SR, McCann L, Fukai T, Holland SM, Mitch WE, Harrison DG. Oxidation of tetrahydrobiopterin leads to uncoupling of endothelial cell nitric oxide synthase in hypertension. *J Clin Invest* 2003;111:1201-9.

Li Y, Zhu H, Kuppasamy P, Roubaud V, Zweier JL, Trush MA. Validation of lucigenin (bis-N-methylacridinium) as a chemiluminescent probe for detecting superoxide anion radical production by enzymatic and cellular systems. *J Biol Chem* 1998;273:2015-23.

Liu Y, Zhao H, Li H, Kalyanaraman B, Nicolosi AC, Gutterman DD. Mitochondrial sources of H₂O₂ generation play a key role in flow-mediated dilation in human coronary resistance arteries. *Circ Res* 2003;93:573-80.

Marchesi E, Rota C, Fann YC, Chignell CF, Mason RP. Photoreduction of the fluorescent dye 2'-7'-dichlorofluorescein: a spin trapping and direct electron spin resonance study with implications for oxidative stress measurements. *Free Radic Biol Med* 1999;26:148-61.

Matés JM, Segura JA, Alonso FJ, Márquez J. Intracellular redox status and oxidative stress: implications for cell proliferation, apoptosis, and carcinogenesis. *Arch Toxicol* 2008;82:271-2.

Pardo Andreu GL, Inada NM, Vercesi AE, Curti C. Uncoupling and oxidative stress in liver mitochondria isolated from rats with acute iron overload. *Arch Toxicol* 2009;83:47-53.

Pryor WA, Houk KN, Foote CS, Fukuto JM, Ignarro LJ, Squadrito GL, Davies KJA. Free radical biology and medicine: it's a gas, man! *Am J Physiol* 2006;291:R491-511.

Rota C, Chignell CF, Mason RP. Evidence for free radical formation during the oxidation of 2'-7'-dichlorofluorescein to the fluorescent dye 2'-7'-dichlorofluorescein by horseradish peroxidase: possible implications for oxidative stress measurements. *Free Radic Biol Med* 1999a;27:873-81.

Rota C, Fann YC, Mason RP. Phenoxyl free radical formation during the oxidation of the fluorescent dye 2',7'-dichlorofluorescein by horseradish peroxidase. Possible consequences for oxidative stress measurements. *J Biol Chem* 1999b;274:28161-8.

Schug M, Heise T, Bauer A, Storm D, Blaszkewicz M, Bedawy E, Brulport M, Geppert B, Hermes M, Föllmann W, Rapp K, Maccoux L, Schormann W, Appel KE, Oberemm A, Gundert-Remy U, Hengstler JG. Primary rat hepatocytes as in vitro system for gene expression studies: comparison of sandwich, Matrigel and 2D cultures. *Arch Toxicol* 2008;82:923-31.

Schumann A, Bauer A, Hermes M, Gilbert M, Hengstler JG, Wilhelm C. A rapid and easy to handle thermoluminescence based technique for evaluation of carbon tetrachloride-induced oxidative stress on rat hepatocytes. *Arch Toxicol* 2009;83:709-20.

- Sebai H, Ben-Attia M, Sani M, Aouani E, Ghanem-Boughanmi N. Protective effect of resveratrol in endotoxemia-induced acute phase response in rats. *Arch Toxicol* 2009;83:335-40.
- Seshiah PN, Weber DS, Rocic P, Valppu L, Taniyama Y, Griendling KK. Angiotensin II stimulation of NAD(P)H oxidase activity: upstream mediators. *Circ Res* 2002;91:406-13.
- Setsukinai K, Urano Y, Kakinuma K, Majima HJ, Nagano T. Development of novel fluorescence probes that can reliably detect reactive oxygen species and distinguish specific species. *J Biol Chem* 2003;278:3170-5.
- Tangkosakul T, Tantimongcolwat T, Isarankura-Na-Ayudhya C, Mejare M, Bülow L, Prachayasittikul V. Native and chimeric metal-binding lactate dehydrogenase as detection and protection tools for oxidative stress induced by Fenton's reaction. *EXCLI J* 2009;8:1-11.
- Tarpey MM, Fridovich I. Methods of detection of vascular reactive species: nitric oxide, superoxide, hydrogen peroxide, and peroxynitrite. *Circ Res* 2001;89:224-36.
- Tarpey MM, Wink DA, Grisham MB. Methods for detection of reactive metabolites of oxygen and nitrogen: in vitro and in vivo considerations. *Am J Physiol* 2004;286:R431-44.
- Thannickal VJ, Fanburg BL. Reactive oxygen species in cell signalling. *Am J Physiol* 2000;279:L1005-28.
- Verstraeten SV, Aimo L, Oteiza PI. Aluminium and lead: molecular mechanisms of brain toxicity. *Arch Toxicol* 2008;82:789-802.
- Wang L, Wang H, Hu M, Cao J, Chen D, Liu Z. Oxidative stress and apoptotic changes in primary cultures of rat proximal tubular cells exposed to lead. *Arch Toxicol* 2009;83:417-27.
- Wrona M, Wardman P. Properties of the radical intermediate obtained on oxidation of 2',7'-dichlorodihydrofluorescein, a probe for oxidative stress. *Free Radic Biol Med* 2006; 41:657-67.
- Yang CS, Lambert JD, Sang S. Antioxidative and anti-carcinogenic activities of tea polyphenols. *Arch Toxicol* 2009;83:11-21.
- Yao DD, Vlessidis AG, Evmiridis NP. Determination of nitric oxide in biological samples. *Microchim Acta* 2004;147:1-20.
- Yu YN, Chen H, Li Y. Protect effect of bicyclol on cisplatin-induced nephrotoxicity in mice. *Arch Toxicol* 2009;83:381-7.
- Zhao H, Kalivendi S, Zhang H, Joseph J, Nithipatikom K, Vásquez-Vivar J, Kalyanaraman B. Superoxide reacts with hydroethidine but forms a fluorescent product that is distinctly different from ethidium: potential implications in intracellular fluorescence detection of superoxide. *Free Radic Biol Med* 2003;34:1359-68.
- Zhao H, Joseph J, Fales HM, Sokoloski EA, Levine RL, Vasquez-Vivar J, Kalyanaraman B. Detection and characterization of the product of hydroethidine and intracellular superoxide by HPLC and limitations of fluorescence. *Proc Natl Acad Sci U S A* 2005;102:5727-32. Erratum in: *Proc Natl Acad Sci U S A* 2005;102:9086.
- Zhou M, Diwu Z, Panchuk-Voloshina N, Haugland RP. A stable nonfluorescent derivative of resorufin for the fluorometric determination of trace hydrogen peroxide: applications in detecting the activity of phagocyte NADPH oxidase and other oxidases. *Anal Biochem* 1997;253:162-8.